

QUASIMINIMALITY IN MIXED TSIRELSON SPACES

ANTONIS MANOUSSAKIS AND ANNA MARIA PELCZAR

ABSTRACT. We prove quasiminimality of the regular mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ with the sequence $(\frac{\theta_n}{\theta_n})_n$ decreasing, where $\theta = \lim_n \theta_n^{1/n}$, and quasiminimality of all mixed Tsirelson spaces $T[(\mathcal{A}_n, \theta_n)_n]$. We prove that under certain assumptions on the sequence $(\theta_n)_n$ the dual spaces are quasiminimal.

INTRODUCTION

Mixed Tsirelson spaces form an important class of spaces in the theory of Banach spaces extensively studied with respect to their distortability, local structure as well as minimality properties.

Recall that an infinite dimensional Banach space X is *minimal*, if any closed infinite dimensional subspace of X contains a further subspace isomorphic to X . A infinite dimensional Banach space X is *quasiminimal*, if any two infinite dimensional subspaces Y, Z of X contains further two infinite dimensional subspaces $Y' \subset Y, Z' \subset Z$ which are isomorphic. The use of Ramsey theory in the famous Gowers' dichotomy inspired studies on minimality and related properties, e.g. [11, 12, 10], in particular in mixed Tsirelson spaces, cf. also [13, 14, 17].

We recall here briefly results concerning minimality properties of mixed Tsirelson spaces. The famous Schlumprecht space $S = T[(\mathcal{A}_n, 1/\log_2(n+1))_n]$ is complementably minimal by [3], i.e. any closed infinite dimensional subspace of S contains a complemented copy of the whole space. This result was proved also for a class of superreflexive spaces extending the construction of S in [9] and for certain types of mixed Tsirelson spaces $T[(\mathcal{A}_{k_n}, \theta_n)_n]$ in [17]. On other hand Tzafriri space $T[(\mathcal{A}_n, \frac{1}{2\sqrt{n}})_n]$ is not minimal [12].

In the original Tsirelson space $T[\mathcal{S}_1, 1/2]$ every normalized block sequence is equivalent to a subsequence of the basis. In context of mixed Tsirelson spaces $T[(\mathcal{S}_{k_n}, \theta_n)_n]$ a more general property - subsequential minimality - was studied. A space with a basis is called *subsequentially minimal*, if any block subspace contains a normalized block sequence equivalent to a subsequence of the basis. In [14] it was proved that if a regular sequence (θ_n) satisfies the condition $(\star)$:

$$\lim_m \limsup_n \frac{\theta_{m+n}}{\theta_n} > 0$$

then the mixed Tsirelson space $X = T[(\mathcal{S}_n, \theta_n)_n]$ is subsequentially minimal if and only if any block subspace of X admits an $\ell_1 - \mathcal{S}_\omega$ -spreading model, if and only if any block subspace of X has Bourgain ℓ_1 -index greater than ω^ω . In particular X is complementably subsequentially minimal if $\sup \theta_n^{1/n} = 1$ [17], i.e. subspaces shown to be isomorphic to subspaces generated by subsequences of the basis are complemented in the space. In [13] analogues of these results were studied in the partly modified setting, in particular it was shown that a partly modified mixed Tsirelson space is subsequentially minimal provided $\sup \theta_n^{1/n} = 1$. Also in [13] a large class of mixed Tsirelson spaces failing the subsequential minimality in a strong sense was described.

In the paper we study p -spaces $T[(\mathcal{A}_n, \theta_n)_n]$, as defined in [17], as well as mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ showing quasiminimality of all spaces of the first type and regular spaces of the second type provided the sequence $(\frac{\theta_n}{\theta})_n$ is decreasing, where $\theta = \lim_n \theta_n^{1/n}$, and we consider related properties. Notice that for the spaces of the second type, in case $\theta < 1$, the monotonicity of the sequence $(\frac{\theta_n}{\theta})_n$ excludes the condition $(\star)$ of [14] mentioned above.

In the study of spaces of both types special averages provide the major tool. In case of p -space we show that ℓ_p is finitely block represented in all block subspaces of any p -spaces, i.e. p is in Krivine set of all block subspaces of a p -space. Equivalent block sequences in these spaces are formed by long averages of ℓ_p -averages.

In case of mixed Tsirelson spaces of type $T[(\mathcal{S}_n, \theta_n)_n]$ we use special averages, whose properties were studied in [2], vectors of the same type as special convex combinations used in [4, 5]. In this case, thanks to the higher complexity of the families $(\mathcal{S}_n)_n$, equivalent sequences are formed by long special averages (not averages of special averages, as it is in p -spaces case). The reasoning proving the quasiminimality of spaces $T[(\mathcal{S}_n, \theta_n)_n]$ implies also the result of [17] on subsequential minimality of the spaces in case $\sup \theta_n^{1/n} = 1$. In both cases the presented argument gives saturation of considered mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ or $T[(\mathcal{A}_n, \theta_n)_n]$ with subsequentially minimal subspaces.

Notice that in both classes of mixed Tsirelson spaces special averages were used in the proofs of arbitrary distortability of spaces under additional assumptions. The observations presented in the paper indicates also similarities in the way spaces ℓ_p , $1 \leq p < \infty$, and Tsirelson type spaces $T[\mathcal{S}_1, \theta]$, $0 < \theta \leq 1$, are represented in spaces $T[(\mathcal{A}_n, \theta_n)_n]$ and $T[(\mathcal{S}_n, \theta_n)_n]$.

Now we describe the contents of the paper. In Section 1 we recall basic notions. In Section 2. we consider p -spaces, state technical lemmas needed in the sequel and describe representation of ℓ_p in p -spaces $T[(\mathcal{A}_n, \theta_n)_n]$ depending on the behavior of $(\theta_n)_n$. The essential tool for quasiminimality in p -spaces case is provided by Theorem 2.8 stating that the Krivine set of any subspace of a p -space, $1 \leq p < \infty$ contains p . Recall here that by [17] Krivine set of a p -space $T[(\mathcal{A}_n, \theta_n)_n]$ contains only p provided the sequence $(\theta_n n^{1/q})_n$ is decreasing, for $1/p + 1/q = 1$, while there are p -spaces, $p > 1$, with 1 in the Krivine set. We present also analogues to observations concerning the original Tsirelson space and mixed Tsirelson spaces given in [19, 20], in particular extending the result of [12] on saturation of Tzafriri space by ℓ_2 -asymptotic subspaces. Section 3 contains the proof of quasiminimality of arbitrary p -spaces (Theorem 3.1) and study of behavior of p -spaces, $1 < p < \infty$, and their duals in case $\inf_n \theta_n n^{1/q} > 0$ (Theorem 3.5), including in particular Tzafriri space. Section 4 is devoted to mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$. After technical lemmas concerning Schreier families, we show the existence of special averages (Corollary 4.9), applying the reasoning of [2] in case the sequence $(\frac{\theta_n}{\theta})_n$ is decreasing. Under this assumption, using special averages we prove the quasiminimality of a regular mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ (Theorem 4.10). Section 5 presents a general argument enabling us to transfer the minimality and quasiminimality properties of spaces to their duals provided certain complementability property holds. In particular we prove quasiminimality of duals to mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ in case $\sup \theta_n^{1/n} = 1$.

1. PRELIMINARIES

We recall now the basic definitions and standard notation. Let X be a Banach space with a basis (e_i) . The *support* of a vector $x = \sum_i x_i e_i$ is the set $\text{supp } x = \{i \in \mathbb{N} : x_i \neq 0\}$, the *range* of a vector $x \in X$ - the smallest interval in $\mathbb{N}$ containing

support of x . Given any $x = \sum_i x_i e_i$ and finite $E \subset \mathbb{N}$ put $Ex = x_E = \sum_{i \in E} x_i a_i$. We write $x < y$ for vectors $x, y \in X$, if $\max \text{supp } x < \min \text{supp } y$. A *block sequence* is any sequence $(x_i) \subset X$ satisfying $x_1 < x_2 < \dots$, a *block subspace* of X - any closed subspace spanned by an infinite block sequence. A subspace spanned by a basic sequence (x_n) we denote by $[x_n]$.

A Banach space X is *saturated* with subspaces of type $(*)$, if any infinite dimensional subspace of X contains a further infinite dimensional subspace of type $(*)$.

A basic sequence (x_n) *C-dominates* a basic sequence (y_n) , $C \geq 1$, if for any scalars (a_n) we have

$$\left\| \sum_n a_n y_n \right\| \leq C \left\| \sum_n a_n x_n \right\|$$

Two basic sequences (x_n) and (y_n) are *C-equivalent*, $C \geq 1$, if (x_n) *C-dominates* (y_n) and (y_n) *C-dominates* (x_n) .

Let $\mathcal{M}$ be a family of finite subsets of $\mathbb{N}$. We say that $\mathcal{M}$ is *regular*, if it is *hereditary*, i.e. for any $G \subset F$, $F \in \mathcal{M}$ also $G \in \mathcal{M}$, *spreading*, i.e. for any integers $n_1 < \dots < n_k$ and $m_1 < \dots < m_k$ with $n_i \leq m_i$, $i = 1, \dots, k$, if $(n_1, \dots, n_k) \in \mathcal{M}$ then also $(m_1, \dots, m_k) \in \mathcal{M}$, and *compact* in the product topology of $2^{\mathbb{N}}$. Given a spreading family $\mathcal{M}$ we say that a sequence $E_1 < \dots < E_n$ of finite subsets of $\mathbb{N}$ is *M-admissible*, if $(\min E_i)_{i=1}^n \in \mathcal{M}$. For any two families $\mathcal{M}, \mathcal{N}$ of finite subsets of $\mathbb{N}$ we define

$$\mathcal{M}[\mathcal{N}] = \{F_1 \cup \dots \cup F_k : F_1, \dots, F_k \in \mathcal{N}, F_1 < \dots < F_k - \mathcal{M} - \text{admissible}\}$$

We will work on two types of regular families: $(\mathcal{A}_n)_{n \in \mathbb{N}}$ and $(\mathcal{S}_\alpha)_{\alpha < \omega_1}$. Let

$$\mathcal{A}_n = \{F \subset \mathbb{N} : \#F \leq n\}, \quad n \in \mathbb{N}$$

Schreier families $(\mathcal{S}_\alpha)_{\alpha < \omega_1}$, introduced in [1], are defined by induction:

$$\mathcal{S}_0 = \{\{n\} : n \in \mathbb{N}\} \cup \{\emptyset\}, \quad \mathcal{S}_{\alpha+1} = \mathcal{S}_1[\mathcal{S}_\alpha], \quad \alpha < \omega_1$$

If α is a limit ordinal, choose $\alpha_n \nearrow \alpha$ and set

$$\mathcal{S}_\alpha = \{F : F \in \mathcal{S}_{\alpha_n} \text{ and } n \leq F \text{ for some } n \in \mathbb{N}\}$$

It is well known that all families $(\mathcal{A}_n)_{n \in \mathbb{N}}$, $(\mathcal{S}_\alpha)_{\alpha < \omega_1}$ are regular.

Let X be a Banach space with a basis. We say that a sequence $x_1 < \dots < x_n$ is *M-admissible*, if $(\text{supp } x_i)_{i=1}^n$ is *M-admissible*.

Definition 1.1 (Tsirelson-type space). *Fix a regular family $\mathcal{M}$ and $0 < \theta < 1$. Let $K \subset c_{00}$ be the smallest set satisfying the following:*

- (1) $(\pm e_n^*)_n \subset K$,
- (2) *for any functionals $\phi_1 < \dots < \phi_k$ in K , if $(\phi_i)_{i=1}^k$ is $\mathcal{M}$ -admissible, then $\theta(\phi_1 + \dots + \phi_k) \in K$.*

We define a norm on c_{00} by the formula $\|x\| = \sup\{f(x) : f \in K\}$, $x \in c_{00}$. Then the Tsirelson-type space $T[\mathcal{M}, \theta]$ is the completion of $(c_{00}, \|\cdot\|)$.

It is standard to verify that the norm $\|\cdot\|$ is the unique norm on c_{00} satisfying the equation

$$\|x\| = \max \left\{ \|x\|_\infty, \sup \left\{ \theta \sum_{i=1}^k \|E_i x\| : (E_i)_{i=1}^k - \mathcal{M} - \text{admissible} \right\} \right\}$$

Recall that $T[\mathcal{A}_n, \theta] = c_0$ if $\theta \leq 1/n$ and $T[\mathcal{A}_n, \theta] = \ell_p$, if $\theta = 1/\sqrt[n]{n}$ for q satisfying $1/p + 1/q = 1$. The space $T[\mathcal{S}_1, 1/2]$ is the Tsirelson space, the first discovered non-classical space - not containing ℓ_p , $1 \leq p < \infty$ or c_0 .

Now we recall the definition of the mixed Tsirelson spaces, using not only one family $\mathcal{M}$ and parameter θ , but sequences of these objects. Again the norm can be defined by its norming set or as a solution of an equation.

Definition 1.2 (Mixed Tsirelson space). *Fix a sequence of regular families $(\mathcal{M}_n)$ and sequence $(\theta_n) \subset (0, 1)$ with $\lim_{n \rightarrow \infty} \theta_n = 0$. Let $K \subset c_{00}$ be the smallest set satisfying the following:*

- (1) $(\pm e_n^*)_n \subset K$,
- (2) for any $\phi_1 < \dots < \phi_k$ in K , if $(\phi_i)_{i=1}^k$ is $\mathcal{M}_n$ -admissible for some $n \in \mathbb{N}$, then $\theta_n(\phi_1 + \dots + \phi_k) \in K$.

We define a norm on c_{00} by $\|x\| = \sup\{f(x) : f \in K\}$, $x \in c_{00}$. The mixed Tsirelson space $T[(\mathcal{M}_n, \theta_n)_n]$ is the completion of $(c_{00}, \|\cdot\|)$.

Again it is standard to verify that the norm $\|\cdot\|$ is the unique norm on c_{00} satisfying the equation

$$\|x\| = \max \left\{ \|x\|_\infty, \sup \left\{ \theta_n \sum_{i=1}^k \|E_i x\| : (E_i)_{i=1}^k \text{ is } \mathcal{M}_n\text{-admissible, } n \in \mathbb{N} \right\} \right\}$$

It follows immediately that the unit vector basis (e_n) is 1-unconditional in the space $T[(\mathcal{M}_n, \theta_n)_n]$. It was proved in [4] that any $T[(\mathcal{S}_{k_n}, \theta_n)_n]$ is reflexive, also any $T[(\mathcal{A}_{k_n}, \theta_n)_n]$ is reflexive, provided $\theta_n > \frac{1}{k_n}$ for at least one $n \in \mathbb{N}$. In this setting Schlumprecht space S is the space $T[(\mathcal{A}_n, \frac{1}{\log_2(n+1)})_n]$, Tzafriri space is $T[(\mathcal{A}_n, \frac{c}{\sqrt{n}})_n]$ for $0 < c < 1$.

A *tree* is a partially ordered set $(T, \prec)$ such that for any $t \in T$ the set $\{s \in T : s \prec t\}$ is finite and linearly ordered. Given a tree T for any $t \in T$ by $\text{succ } t$ we denote the set of immediate successors of t in T . The *height* of a tree T is the supremum of the length of branches in the tree T (i.e. well ordered subsets of the tree T). The *level* of an element t of a tree T with a unique root (i.e. the minimal element) is the length of the branch linking the root with t .

The following notion provides a useful tool for estimating norms in mixed Tsirelson spaces:

Definition 1.3 (The tree-analysis of a norming functional). *Let $\phi \in K$. By a tree-analysis of ϕ we mean a finite family $(\phi_t)_{t \in T}$ indexed by a tree T with a unique root $0 \in T$ such that the following hold*

- (1) $\phi_0 = \phi$ and $\phi_t \in K$ for all $t \in T$,
- (2) $t \in T$ is maximal if and only if $\phi_t \in (\pm e_n^*)_n$,
- (3) for every $t \in T$ not maximal there is some $n \in \mathbb{N}$ such that $(\phi_s)_{s \in \text{succ } t}$ is an $\mathcal{M}_n$ -admissible block sequence and $\phi_t = \theta_n(\sum_{s \in \text{succ } t} \phi_s)$. We call θ_n the weight of ϕ_t .

Notice that every functional from a norming set K admits a tree-analysis, not necessarily unique.

Definition 1.4. *Let $(y_n) \subset X$ be a block sequence.*

- a) *Let $\phi \in K$ be a norming functional with a tree-analysis $(\phi_t)_{t \in T}$. We say that ϕ_t covers y_n if $t \in T$ is maximal in T with the property $\text{supp } \phi_t \supset \text{supp } y_n \cap \text{supp } \phi$.*
- b) *a finite sequence $(E_k)_{k=1}^m$ of subsets of $\mathbb{N}$ is said to be comparable with the sequence (y_n) if for every n and every $k \leq m$ we have*

$$E_k \subset \text{range } y_n \text{ or } \text{supp } y_n \cap \bigcup_{i=1}^m E_i \subset E_k \text{ or } \text{range } E_k \cap \text{range } y_n = \emptyset.$$

- c) *A norming functional $\phi \in K$ with a tree-analysis $(\phi_t)_{t \in T}$ is said to be comparable with $(y_n)_n$ if the sequence $(\text{supp } \phi_t)_{t \in T}$ is comparable with $(y_n)_n$.*

2. REPRESENTABILITY OF ℓ_p IN p -SPACES

We describe now the class of p -spaces following the definition of [17]. Recall that any space $T[(\mathcal{A}_{k_n}, \theta_n)_{n=1}^N]$, defined analogously to Tsirelson type spaces and mixed Tsirelson spaces, with the use of finitely many families $(\mathcal{A}_{k_n})_{n=1}^N$ and finitely many scalars $(\theta_n)_{n=1}^N \subset (0, 1)$, is isomorphic to some ℓ_p , $1 \leq p < \infty$ or c_0 [7].

Definition 2.1. [17] *A mixed Tsirelson space $T[(\mathcal{A}_{k_n}, \theta_n)_{n \in \mathbb{N}}]$ is called a p -space, for $p \in [1, \infty)$, if there is a sequence $(p_N)_N \subset (1, \infty)$ such that*

- (1) $p_N \rightarrow p$ as $N \rightarrow \infty$, and $p_N \geq p_{N+1} > p$ for any $N \in \mathbb{N}$,
- (2) $T[(\mathcal{A}_{k_n}, \theta_n)_{n=1}^N]$ is isomorphic to ℓ_{p_N} for any $N \in \mathbb{N}$.

A p -space $T[(\mathcal{A}_n, \theta_n)_{n \in \mathbb{N}}]$ is called regular, if $\theta_n \searrow 0$ and $\theta_{nm} \geq \theta_n \theta_m$ for any $n, m \in \mathbb{N}$.

Notice that any mixed Tsirelson space $T[(\mathcal{A}_{k_n}, \theta_n)_{n \in \mathbb{N}}]$ is isometric to a space $T[(\mathcal{A}_n, \hat{\theta}_n)_{n \in \mathbb{N}}]$, where

$$\hat{\theta}_n = \sup \left\{ \prod_{i=1}^l \theta_{n_i} : \prod_{i=1}^l n_i \geq n \right\}, \quad n \in \mathbb{N}$$

It follows directly that if $T[(\mathcal{A}_{k_n}, \theta_n)_{n \in \mathbb{N}}]$ is a p -space, then also $T[(\mathcal{A}_n, \hat{\theta}_n)_{n \in \mathbb{N}}]$ is a p -space. Notice also that the sequence $(\hat{\theta}_n)$ satisfies $\hat{\theta}_n \searrow 0$, $\hat{\theta}_{nm} \geq \hat{\theta}_n \hat{\theta}_m$ for any $n, m \in \mathbb{N}$.

Therefore from now on we assume that p -spaces we consider are regular.

Notation. Let $T[(\mathcal{A}_n, \theta_n)_{n \in \mathbb{N}}]$ be a regular p -space. If we set $\theta_n = 1/n^{1/q_n}$ with $q_n \in (1, \infty)$, $n \in \mathbb{N}$, then $q = \lim_n q_n = \sup_n q_n \in (0, \infty]$, where $1/p + 1/q = 1$, with usual convention $1/\infty = 0$.

In the situation as above let $c_n = \theta_n n^{1/q} \in (0, 1)$, $n \in \mathbb{N}$, if $p > 1$. To unify the notation put $c_n = \theta_n$, $n \in \mathbb{N}$, in case $p = 1$.

Remark 2.2. In [17] in the definition in the p -spaces it was posed also that the sequence $(p_n)_n$ was strictly decreasing and hence θ_n also. It was also posed that $c_n \searrow 0$. In our setting, the sequences $(p_n)_n, (c_n)_n$ do not necessarily satisfy these two conditions.

Remark 2.3. Notice that the norming set of a p -space is contained in the unit ball of ℓ_q , hence $\|\cdot\| \leq \|\cdot\|_p$.

First we present two technical observations.

Lemma 2.4. *Let X be a p -space, $1 \leq p < \infty$. Take a functional ϕ with a tree-analysis $(\phi_t)_{t \in T}$ and any finite block sequence (v_n) . Set $v = \sum_n v_n$. Then there is a functional ϕ' with a tree-analysis $(\phi'_t)_{t \in T'}$ comparable with (v_n) satisfying $6\phi'(v) \geq \phi(v)$.*

Proof. We can assume that $\text{supp } \phi \subset \bigcup_n \text{supp } v_n$. For any n pick $t_n \in T$ so that ϕ_{t_n} covers v_n . Let $E_1 < \dots < E_k$ be supports of all successors $\phi_1, \dots, \phi_k$ of ϕ_{t_n} which intersect the support of v_n .

CASE 1. There is $i = 1, \dots, k$ so that $E_i \subset \text{supp } v_n$. If $E_1 \cap \text{supp } v_{n-1} \neq \emptyset$ and $\phi_1(v_n) \geq \phi_i(v_n)$ then change the tree: split ϕ_1 into two parts supported on $E_1 \cap \text{supp } v_n$ and $E_1 \setminus \text{supp } v_n$ and erase ϕ_i , put the part $\phi_1|_{\text{supp } v_n}$ into tree instead of ϕ_i , leave $\phi_1|_{E_1 \setminus \text{supp } v_n}$ in the place of ϕ_1 . If $E_1 \cap \text{supp } v_{n-1} \neq \emptyset$ and $\phi_1(v_n) \leq \phi_i(v_n)$ then erase from ϕ the part supported on $E_1 \cap \text{supp } v_n$. Analogously proceed, if $E_k \cap \text{supp } v_{n+1} \neq \emptyset$. The action of the modified functional on v_n is not less than $\phi(v_n)/3$.

CASE 2. $k = 2$, $E_1 \cap \text{supp } v_{n-1} \neq \emptyset$, $E_2 \cap \text{supp } v_{n+1} \neq \emptyset$. Let for example $\phi_1(v_n) \geq \phi_2(v_n)$, then erase part of ϕ supported on $E_2 \cap \text{supp } v_n$, and for the modified functional find new t_n - then we have only Case 1. $\square$

Lemma 2.5. *Let X be a p -space, $1 \leq p < \infty$. Take any norming functional ϕ with a tree-analysis $(\phi_t)_{t \in T}$ and some $J_1, \dots, J_N \subset T$ so that*

- (1) $J_n \subset \text{succ } t_n$ for some $t_n \in T$ for any $n = 1, \dots, N$,
- (2) $\text{supp } \phi_i \cap \text{supp } \phi_j = \emptyset$ for any $i \neq j$, $i, j \in \bigcup_n J_n$
- (3) $\text{supp } \phi = \bigcup \{\text{supp } \phi_i : i \in \bigcup_n J_n\}$

Let $\phi = \sum_{n=1}^N \gamma_n \sum_{i \in J_n} \phi_i$. Then for any scalars $a_1, \dots, a_N \geq 0$ we have

$$\sum_{n=1}^N a_n \gamma_n (\#J_n)^{1/q} \leq \left(\sum_{n=1}^N a_n^p \right)^{1/p}$$

Proof. The case $p = 1$ is obvious. Assume $p > 1$. We proceed by induction on the height of the tree T . For the height equal 0 (i.e. ϕ is a unit vector) the result is clear. Assume we have the result for functionals with the tree-analysis of height smaller than l for some $l \geq 1$ and pick a functional with the tree-analysis $(\phi_t)_{t \in T}$ of height l . Let $J_1, \dots, J_N, \gamma_1, \dots, \gamma_N, a_1, \dots, a_N$ be as in the Lemma.

Let $\phi = c_m / m^{1/q} \sum_{t \in \text{succ } 0} \phi_t$. Let $I = \{n = 1, \dots, N : J_n \subset \text{succ } 0\}$. Let $J = \text{succ } 0 \setminus \bigcup_n J_n$. For any $t \in J$ let $I_t = \{n \notin I : i \succ t \ \forall i \in J_n\}$. By the inductive assumption for ϕ_t , $t \in J$, and Hölder inequality we have

$$\begin{aligned} \sum_{n=1}^N \gamma_n a_n (\#J_n)^{1/q} &\leq \frac{1}{m^{1/q}} \sum_{n \in I} a_n (\#J_n)^{1/q} + \frac{1}{m^{1/q}} \sum_{t \in J} \sum_{n \in I_t} a_n \gamma_n m^{1/q} (\#J_n)^{1/q} \\ &\leq \frac{1}{m^{1/q}} \left(\sum_{n \in I} a_n^p \right)^{1/p} \left(\sum_{n \in I} \#J_n \right)^{1/q} + \frac{1}{m^{1/q}} \sum_{t \in J} \left(\sum_{n \in I_t} a_n^p \right)^{1/p} \\ &\leq \frac{1}{m^{1/q}} \left(\sum_{n \in I} a_n^p \right)^{1/p} \left(\sum_{n \in I} \#J_n \right)^{1/q} + \frac{1}{m^{1/q}} (\#J)^{1/q} \left(\sum_{t \in J} \sum_{n \in I_t} a_n^p \right)^{1/p} \end{aligned}$$

Notice that $I \cup \bigcup_{t \in J} I_t = \{1, \dots, N\}$ and $\#J + \sum_{n \in I} \#J_n = \# \text{succ } 0 \leq m$, hence again by Hölder inequality we obtain the desired upper bound by $(\sum_{n=1}^N a_n^p)^{1/p}$. $\square$

We recall here the notion of ℓ_r -averages - the basic tool we will use in studying the properties of p -spaces:

Definition 2.6. *A vector $x \in X$ is called a $C - \ell_r$ -average of length m , for $r \in [1, \infty]$, $m \in \mathbb{N}$ and $C \geq 1$ if $x = \sum_{n=1}^m x_i / \|\sum_{n=1}^m x_i\|$ for some normalized block sequence $(x_n)_{n=1}^m$ which is C -equivalent to the unit vector basis of ℓ_r^m .*

The following lemma extends the standard result for ℓ_1 -averages.

Lemma 2.7. *Fix $1 \leq r < \infty$ and $M \in \mathbb{N}$. Let $N \geq (2M)^r$. Then for any $C - \ell_r$ -average $x \in X$ of length N and any $j \leq M$ we have*

$$\frac{j^{1/s}}{2C^2} \leq \sup \left\{ \sum_i \|E_i x\| : (E_i) - \mathcal{A}_j - \text{admissible} \right\} \leq 2C^2 j^{1/s}$$

where $1/s + 1/r = 1$.

Proof. Let $x = (x_1 + \dots + x_N) / \|x_1 + \dots + x_N\|$, for some normalized sequence $x_1 < \dots < x_N$ C -equivalent to the unit vector basis of ℓ_p^N . Fix $j \leq M$. For the lower estimate take subsets $I_1 < \dots < I_j$ of $\{1, \dots, N\}$ such that $\#I_i = \lfloor N/j \rfloor$, i.e.

the integer part of N/j , let $E_i = \bigcup_{n \in I_i} \text{supp } x_n$ and compute

$$\sum_{i=1}^j \|E_i x\| \geq \sum_i \left\| \sum_{n \in I_i} x_n \right\| / C N^{1/r} \geq \frac{j}{C^2 N^{1/r}} \left[\frac{N}{j} \right]^{1/r} \geq \frac{j^{1/s}}{2^{1/r} C^2} \geq \frac{j^{1/s}}{2 C^2}.$$

Now take any $E_1 < \dots < E_j$ for $j \leq M$. For any $i = 1, \dots, j$ pick $I_i = \{n : \text{supp } x_n \cap E_i \neq \emptyset\}$ and $I'_i = \{n \in I_i : \text{supp } x_n \cap E_l = \emptyset \ \forall i \neq l\}$ and compute

$$\|E_i(x_1 + \dots + x_N)\| \leq \|x_{\min I_i}\| + \|x_{\max I_i}\| + \|E_i(\sum_{n \in I'_i} x_n)\| \leq 2 + C(\#I'_i)^{1/r}$$

Hence summing over i we get

$$\sum_i \|E_i x\| \leq 2M \frac{C}{N^{1/r}} + C^2 \sum_i \frac{(\#I'_i)^{1/r}}{N^{1/r}} \leq 2C^2 j^{1/s}$$

□

Now we study the way ℓ_p is represented in a p -space. Recall that by Krivine theorem for any Banach space X with a basis there is some $1 \leq p \leq \infty$ such that ℓ_p is finitely block (almost isometrically) represented in X , i.e. for any $\varepsilon > 0$ and any $n \in \mathbb{N}$ there is a normalized block sequence $x_1 < \dots < x_n$ in X which is $(1 + \varepsilon)$ -equivalent to the unit vector basis of ℓ_p^n . The set of all such p 's is called the Krivine set of a given space X . Recall that the Krivine set of a p -space $T[(\mathcal{A}_n, \theta_n)_{n \in \mathbb{N}}]$ contains only p provided the sequence $(c_n)_n$ is decreasing (Prop. 1.6 [17]), while there are p -spaces with $p > 1$ and 1 in the Krivine set (Prop. 1.8 [17]).

Theorem 2.8. *Let X be a p -space, $1 \leq p < \infty$. Then the Krivine set of any block subspace of X contains p .*

Proof. By Lemma 1.5 [17] Krivine set of X is contained in $[1, p]$. We will show that for any N in any block subspace Y of X there is a normalized block sequence $y_1, \dots, y_N$ with $\|\sum_{n \in J} y_n\| \leq D(\#J)^{1/p}$ for any $J \subset \{1, \dots, N\}$ and for a universal constant D . Then by Corollary 5 [18] some $p' \geq p$ is in the Krivine set of Y , thus $p' = p$ which ends the proof.

Fix $N \in \mathbb{N}$ and $r \leq p$ in the Krivine set of a block subspace Y . Take a normalized block sequence $(y_n)_1^N$ and sequence $(m_n)_1^N \subset \mathbb{N}$ with

- (1) y_n is a $2 - \ell_r$ -average of length greater than $(2m_n)^r$, for any n ,
- (2) $N\theta_{m_n} \leq 1/2^{n+2}$, for any n ,
- (3) $\theta_{m_n} \sum_{i < n} \#\text{supp } y_i \leq 1/2^{n+2}$ for any n ,

We claim that $\|\sum_{n=1}^N y_n\| \leq 99N^{1/p}$.

Take a functional ϕ with the tree analysis $(\phi_t)_{t \in T}$. By Lemma 2.4 we can assume that this tree-analysis is comparable with (y_n) . Let $y = \sum_n y_n$.

Let $A = \{t \in T : \phi_t \text{ covers some } y_n\}$. Given any $t \in A$ let $I_t = \{n = 1, \dots, N : \phi_t \text{ covers } y_n\}$. Let θ_m be the weight of ϕ_t .

Now consider cases:

CASE 1. $m \leq m_n$ for all $n \in I_t$.

For any $n \in I_t$ let $J_n = \{i \in \text{succ } t : \text{supp } \phi_i \subset \text{range } y_n\}$. By Lemma 2.7 we have $\sum_{i \in J_n} \phi_i(y) = \sum_{i \in J_n} \phi_i(y_n) \leq 8(\#J_n)^{1/q}$.

CASE 2. There is some $n \in I_t$ with $m > m_n$. Let n_t be the maximal element of I_t with this property.

First let $L_t = \{n \notin I_t : \text{supp } y_n \cap \text{supp } \phi \subset \text{supp } \phi_t\}$. Notice that for any $n \in L_t$ there is some ϕ_{t_n} -successor of ϕ_t so that $\text{supp } y_n \cap \text{supp } \phi \subset \text{supp } \phi_{t_n}$. Hence

$$\phi_t \left(\sum_{n \in L_t} y_n \right) \leq \theta_{m_{n_t}} \left(\sum_{n \in L_t} \phi_{t_n}(y_n) \right) \leq N \theta_{m_{n_t}} \leq 1/2^{n_t+2}$$

Hence $\phi(\sum_{t \in A, n \in L_t} y_n) \leq 1/4$, thus we can erase this part for all t with error $1/4$.

Now notice that by condition 3. we have

$$\phi_t \left(\sum_{n \in I_t, n < n_t} y_n \right) \leq \theta_{m_{n_t}} \sum_{n < n_t} \# \text{supp } y_n \leq 1/2^{n_t+2}$$

so we can again erase this part for all t with error $1/4$.

Now we compare the parts $\phi_t(y_{n_t})$ and $\phi_t(\sum_{n \in I_t, n > n_t} y_n)$. If the first one is bigger, then set $J_{n_t} = \{t\}$. Obviously we have $\phi_t(y) \leq (\#J_{n_t})^{1/q}$. If the second one is bigger, then erase the first one and proceed as in Case 1 getting $\sum_{i \in J_n} \phi_i(y) \leq 8(\#J_n)^{1/q}$ for each $n \in I_t, n > n_t$.

Now by Lemma 2.5 we have $\phi(\sum_{n=1}^N y_n) \leq 2 \cdot 8N^{1/p} + \frac{1}{2}$, therefore $\|\sum_{n=1}^N y_n\| \leq 6 \cdot 16^{\frac{1}{2}} N^{1/p} = 99N^{1/p}$ which ends the proof of Theorem. $\square$

Now we work under more restrictive assumptions on (c_n) , giving additional information on representability of ℓ_p in p -spaces. Recall that a space with a basis is called ℓ_p -asymptotic provided any normalized block sequence $n \leq x_1 < \dots < x_n$, $n \in \mathbb{N}$, is C -equivalent to the unit vector basis of ℓ_p^n , with some universal constant C . Now we show a result generalizing Corollary 3.8 [12].

Proposition 2.9. *Let X be a p -space, $1 < p < \infty$, with $\inf_n c_n = c > 0$. Then X is saturated with ℓ_p -asymptotic subspaces.*

To prove the Proposition it is enough to show the following Lemma (well-known in case of Tzafriri space) similar to Proposition 1.6 [17].

Lemma 2.10. *Assume $\inf_n c_n = c > 0$. Take a normalized block sequence $y_1 < \dots < y_N$ in X . Then for any $(a_n)_{n=1}^N$ we have $\|\sum_n a_n y_n\| \leq 6c^{-1}(\sum_n a_n^p)^{1/p}$.*

Proof. Take any norming functional ϕ with the tree-analysis $(\phi_t)_{t \in T}$, by Lemma 2.4 we can assume that the tree-analysis is comparable with (y_n) . For any n pick $t_n \in T$ such that ϕ_{t_n} covers y_n and let $(\phi_i)_{i \in J_n}$ be the family of immediate successors of ϕ_{t_n} with support contained in $\text{supp } y_n$. By assumption and definition of the norm we have $\sum_{i \in J_n} \phi_i(y_n) \leq c^{-1}(\#J_n)^{1/q}$ for any n . Now we apply Lemma 2.5 obtaining $\phi(\sum_n a_n y_n) \leq 6c^{-1}(\sum_n a_n^p)^{1/p}$. $\square$

By this Lemma and the fact that for any normalized $y_1 < \dots < y_N$ also $\|\sum_n y_n\| \geq cN^{1/p}$, applying Theorem 3.7 [12], we obtain saturation of X by ℓ_p -asymptotic subspaces.

Recall that by proof of Theorem 2.1 [17], if $c_n \searrow 0$, then X does not contain a ℓ_p -asymptotic subspace.

The next two observations should be compared with Proposition 3.9 [19] and Example 5.12 [20], the idea of the proofs is analogous.

Proposition 2.11. *Let X be a p -space, $1 < p < \infty$, with $\prod_n c_n = d > 0$ or $c_n \nearrow 1$. Then X is saturated with ℓ_p .*

Proof. By quasiminimality (Theorem 3.1) it is enough to show that there is a subspace isomorphic to ℓ_p . Since we have for any normalized block sequence the upper ℓ_p -estimate by Lemma 2.10, it is enough to produce a normalized block sequence with lower ℓ_p -estimate. We present the proof in case $\prod_n c_n = d > 0$, the reasoning can be easily adapted to the case $c_n \nearrow 1$.

Take increasing $(i_n)_n \subset \mathbb{N}$ with $i_{n+1} > 5i_n$ and $\sum_n 2^{-i_n} \leq d/4$, and take the partition $I_1 < I_2 < \dots$ of $\mathbb{N}$ with $\#I_n = 2^{i_n}$. Notice that by Remark 2.3 we have $\|\sum_{i \in I_n} e_i\| \leq 2^{i_n/p}$.

Let $x_n = (\|\sum_{i \in I_n} e_i\|)^{-1} \sum_{i \in I_n} e_i$ for any n . We claim that (x_n) satisfies the lower ℓ_p -estimate.

Take $(j_n)_{n=1}^N \subset \mathbb{N}$ with $j_n \leq i_n$ and consider the vector $x = \sum_n 2^{-j_n/p} x_n$. Let $k_n = j_n + i_n$ for any n , then $k_{n+1} > k_n$. Then the formula

$$\phi = \sum_{n=1}^N \sum_{i \in I_n} (c_{2^{k_1}} c_{2^{k_2-k_1}} \dots c_{2^{k_n-k_{n-1}}}) 2^{-k_n/q} e_i$$

defines a functional from the norming set with a tree-analysis all of whose elements have weights in the set $\{\theta_{2^{k_1}}, \theta_{2^{k_2-k_1}}, \dots, \theta_{2^{k_n-k_{n-1}}} : n = 1, \dots, N\}$. Moreover $k_{n+1} - k_n > k_n - k_{n-1}$ for any n , hence

$$\|x\| \geq \phi(x) \geq \sum_n \left(\prod_k c_k \right) \left(2^{-k_n/q} 2^{-j_n/p} \#I_n / \left\| \sum_{i \in I_n} e_i \right\| \right) \geq d \sum_n 2^{-j_n}$$

For any scalars $(a_n)_{n=1}^N \subset [0, 1]$ with $\sum_n a_n^p = 1$ pick (j_n) with $2^{(-j_n-1)/p} \leq a_n \leq 2^{-j_n/p}$. Let $J = \{n \in \mathbb{N} : j_n \geq i_n\}$. Then by the choice of i_n we have

$$\sum_{n \in J} a_n^p \leq \sum_{n \in J} 2^{-j_n} \leq \sum_n 2^{-i_n} \leq d/4$$

By the previous argument we have

$$\left\| \sum_n a_n x_n \right\| \geq \left\| \sum_{n \notin J} a_n x_n \right\| \geq d \sum_{n \notin J} 2^{-j_n-1} \geq d \sum_n a_n^p / 2 - d/4 = d/4$$

Hence we proved the lower ℓ_p -estimate for (x_n) and thus the proposition. $\square$

Proposition 2.12. *Let X be a p -space, $1 \leq p < \infty$, with $\sup_n c_n = d < 1$. Then X does not contain ℓ_p .*

Proof. Given a normalized block basis (x_j) and $k \in \mathbb{N}$ define a new norm $\|\cdot\|_{k, (x_j)}$ on $[x_j]$ in the following way: for any $x \in [x_j]$ let $\|x\|_{k, (x_j)}$ be the supremum of $\phi(x)$ over all norming functionals ϕ with a tree-analysis $(\phi_t)_{t \in T}$ such that for any $t \in T$ of level $\leq k$ and any j either $\text{supp } \phi_t \cap \text{supp } x_j = \emptyset$ or $\text{range } \phi_t \supset \text{supp } x_j$.

Lemma 2.13. *Let (x_j) be a normalized block basis in X , fix $\varepsilon > 0$ and $k \in \mathbb{N}$. Then there is $x \in [x_j]$ with $\|x\| = 1$ such that $\|x\|_{k, (x_j)} > 1 - \varepsilon$.*

Proof of Lemma. We will prove it first for $k = 1$. Take a normalized block sequence $(y_n^1)_{n=1}^N$ of (x_j) such that

- (1) $y_n^1 = \sum_{j \in F_n} x_j / \|\sum_{j \in F_n} x_j\|$, where $\#F_n = m_n$ for any n ,
- (2) $m_n \sum_{i > n} 1/\theta_{m_i} m_i \leq \varepsilon/4$ for any n ,
- (3) $\theta_{m_n} \sum_{i < n} \#\text{supp } y_i^1 \leq \varepsilon/4$ for any n ,
- (4) $1/N\theta_N \leq \varepsilon/8$.

Let $y^2 = \sum_{n=1}^N y_n^1 / \|\sum_{n=1}^N y_n^1\|$. Take $(E_i)_{i=1}^l$ with $\|y^2\| = \theta_l \sum_i \|E_i y^2\|$. We claim that with error at most ε we can assume that for any n there is at most one i with $\text{supp } x_n \cap E_i \neq \emptyset$. Let $J_n = \{j \in F_n : \text{supp } x_j \text{ intersects at least two } E_i\}$.

Assume that there is some n_0 such that $m_{n_0} < l \leq m_{n_0+1}$. Then we let w_1 be y^2 restricted to supports of y_n^1 for $n < n_0$, w_2 be y^2 restricted to supports of $y_{n_0}^1$ and $y_{n_0+1}^1$. Then $\theta_l \sum_i \|E_i w_1\| \leq \varepsilon/4$ by condition 3., $\theta_l \sum_i \|E_i w_2\| \leq 2/N\theta_N \leq \varepsilon/4$ by condition 4., and by condition 1. we have

$$\sum_{n > n_0+1} \sum_{j \in J_n} \|y|_{\text{supp } x_j}\| \leq m_{n_0+1} \sum_{n > n_0+1} 1/\theta_{m_n} m_n \leq \varepsilon/4$$

Any other case is a simpler modification of the reasoning above.

For any $k > 1$ we iterate the procedure above, by taking as (y_n^j) suitable averages of (y_n^{j-1}) for some ε_j , $j = 2, \dots, k$, with $\prod_{i=1}^k (1 - \varepsilon_i) > 1 - \varepsilon$. $\square$

Now we go back to the proof of the Proposition. Assume there is a normalized block basis (x_j) D -equivalent to the unit vector basis of ℓ_p . Take $k \in \mathbb{N}$ and by the Lemma 2.13 pick a normalized $x = \sum_j a_j x_j$ with $\|x\|_{k, (x_j)} > 1/2$. Let

$$\|x\|_{k, (x_j)} = \sum_i \frac{c_{n_1^i} c_{n_2^i} \dots c_{n_{k_i}^i}}{(n_1^i \dots n_{k_i}^i)^{1/q}} \|E_i x\|$$

for some E_i such that the sets $J_i = \{j : \text{supp } x_j \cap E_i \neq \emptyset\}$ are pairwise disjoint and some $k_i \leq k$ and $(n_1^i, n_2^i, \dots, n_{k_i}^i)_i$ with $\sum_i 1/(n_1^i \dots n_{k_i}^i) \leq 1$. Then by Hölder inequality

$$1/2 \leq d^k \sum_i 1/(n_1^i \dots n_{k_i}^i)^{1/q} D(\sum_{j \in J_i} a_j^p)^{1/p} \leq d^k D(\sum_j a_j^p)^{1/p} \leq d^k D^2$$

which for large k gives contradiction. $\square$

3. QUASIMINIMALITY IN p -SPACES

In this section we prove the following

Theorem 3.1. *Any p -space X , $1 \leq p < \infty$, is quasiminimal.*

We show also a stronger quasiminimality property of X and its dual space under additional assumption on (c_n) . The equivalent sequences used in the proof of Theorem 3.1 consist of averages of ℓ_p -averages, whose properties are examined in next Lemmas.

Lemma 3.2. *Fix $N \in \mathbb{N}$. Take finite normalized block sequences $(y_n)_1^N$ and $(z_n)_1^N$ in X and sequence $(m_n)_1^N \subset \mathbb{N}$ satisfying the following conditions:*

- (1) *each y_n and z_n are $2 - \ell_p$ -averages of length $N_n \geq (2m_n)^p$,*
- (2) *$N\theta_{m_n} \leq 1/2^{n+5}$ for any n ,*
- (3) *$\theta_{m_n} \sum_{i < n} \#\text{supp } y_i \leq 1/2^{n+5}$, and $\theta_{m_n} \sum_{i < n} \#\text{supp } z_i \leq 1/2^{n+5}$ for any n ,*

Then $(y_n)_1^N$ and $(z_n)_1^N$ are C -equivalent, for some universal constant C .

Proof. We show that $(z_n)_1^N$ dominates $(y_n)_1^N$. Fix scalars $0 \leq a_1, \dots, a_N \leq 1$ so that the vector $y = \sum_{n=1}^N a_n y_n$ has norm 1, and put $z = \sum_{n=1}^N a_n z_n$. The proof resembles the reasoning in the proof of Theorem 2.8.

By Lemma 2.4 we can take a norming functional ϕ with a tree analysis $(\phi_t)_{t \in T}$ comparable with (y_n) and $\phi(y) \geq 1/6$. We will pick ψ so that $\|\psi\| \leq 1$ and $\psi(z) \geq 2^{-11}$.

Let $A = \{t \in T : \phi_t \text{ covers some } y_n\}$. Given any $t \in A$ let $I_t = \{n = 1, \dots, N : \phi_t \text{ covers } y_n\}$. Let θ_m be the weight of ϕ_t .

Now consider cases for each $t \in A$:

CASE 1. $m \leq m_n$ for all $n \in I_t$. For any $n \in I_t$ let $J_n = \{i \in \text{succ } t : \text{supp } \phi_i \subset \text{range } y_n\}$. By condition 1. and Lemma 2.7 pick functionals $(\psi_i)_{i \in K_n}$, with $\#K_n \leq \#J_n$ so that $\sum_{i \in K_n} \psi_i(z_n) \geq \sum_{i \in J_n} \phi_i(y_n)2^{-6}$ and change the tree-analysis of ϕ by putting $(\psi_i)_{i \in K_n}$ instead of $(\phi_i)_{i \in J_n}$ for any $i = 1, \dots, k_n$. Then we have $2^6 \psi_t(\sum_{n \in I_t} a_n z_n) \geq \phi_t(\sum_{n \in I_t} a_n y_n)$.

CASE 2. there is some $n \in I_t$ with $m > m_n$. Let n_t be maximal with this property.

First let $L_t = \{n \notin I_t : \text{supp } y_n \cap \text{supp } \phi \subset \text{supp } \phi_t\}$. Notice that for any $n \in L_t$ there is some ϕ_{t_n} - a successor of ϕ_t so that $\text{supp } y_n \cap \text{supp } \phi \subset \text{supp } \phi_{t_n}$. Hence

$$\phi_t(\sum_{n \in L_t} a_n y_n) \leq \theta_{m_{n_t}}(\sum_{n \in L_t} \phi_{t_n}(y_n)) \leq N \theta_{m_{n_t}} \leq 1/2^{n_t+5}$$

Hence $\phi(\sum_{n \in L_t, t \in B} a_n y_n) \leq 1/2^5$, where $B = \{t \in A : m > m_k \text{ for some } k \in I_t\}$, thus we can erase this part with error $1/2^5$.

Now notice that by condition 2. we have

$$\phi_t(\sum_{n \in I_t, n < n_t} a_n y_n) \leq \theta_{m_{n_t}} \sum_{n < n_t} \# \text{supp } y_n \leq 1/2^{n_t+5}$$

so we can again erase this part for all t with error $1/2^5$.

Now we compare the part $\phi_t(a_{n_t} y_{n_t})$ and $\phi_t(\sum_{n \in I_t, n > n_t} a_n y_n)$. If the first one is bigger, then replace ϕ_t in the tree-analysis of ϕ by ψ_t satisfying $\psi_t(z_{n_t}) = 1$. If the second one is bigger, then erase the first one and proceed as in Case 1.

Altogether after all this changes we obtained a new norming functional ψ with $\psi(\sum_n a_n z_n) \geq 2^{-11}$. Indeed, from the above inequalities we have

$$(3.1) \quad \frac{1}{6} \leq \phi(\sum_n y_n) \leq \frac{2}{2^5} + 2 \cdot 2^6 \psi(\sum_n a_n y_n) \Rightarrow \psi(\sum_n a_n y_n) \geq (\frac{1}{6} - \frac{1}{2^4}) \frac{1}{2^7}.$$

Therefore $C = 2^{11}$ satisfies the assertion of the Lemma. $\square$

Lemma 3.3. Fix $N \in \mathbb{N}$. Take $(y_n)_1^N, (z_n)_1^N$ as in Lemma 3.2 and let $y = y_1 + \dots + y_N, z = z_1 + \dots + z_N$. Then for $m \in \mathbb{N}$, any $E_1 < \dots < E_m$ there are $F_1 < \dots < F_l$ with $l \leq m$

$$(3.2) \quad \theta_m(\|E_1 y\| + \dots + \|E_m y\|) \leq C' \theta_m(\|F_1 z\| + \dots + \|F_l z\|) + 6$$

for some universal constant C' .

Proof. We can assume (up to multiplying by 3) that the sequence (E_j) are comparable with the sequence (y_n) . Let $J = \{j = 1, \dots, m : \text{range } E_j \supset \text{supp } y_n \text{ for some } n\}$ and for $j \in J$ let $I_j = \{n : \text{supp } y_n \cap \bigcup_i E_i \subset E_j\}$. Then for any $j \in J$ we put $F_j = \bigcup_{n \in I_j} \text{supp } z_n$ and we have by Lemma 3.2

$$\sum_{j \in J} \|E_j \sum_{n \in I_j} y_n\| \leq C \sum_{j \in J} \|F_j \sum_{n \in I_j} z_n\|$$

Symmetrically let $L = \{n : \text{range } y_n \supset E_j \text{ for some } j\}$ and for any $n \in L$ let $J_n = \{j : E_j \subset \text{range } y_n\}$. Consider cases:

CASE 1. $m \leq m_n$ for any n . Then by the choice of $(y_n), (z_n)$ and Lemma 2.7 there are some $(F_j)_{j \in K_n, n \in I}$, $\#K_n \leq \#J_n$ with

$$\sum_{n \in L} \sum_{j \in J_n} \|E_j y_n\| \leq 2^6 \sum_{n \in L} \sum_{j \in K_n} \|F_j z_n\|$$

and $\{F_j\}_{j \in J} \cup \{F_j\}_{j \in K_n, n \in L}$ are the desired sets.

CASE 2. For some n we have $m > m_n$. Let n_0 be maximal with this property. Then

$$\theta_m \sum_{n \in L, n < n_0} \sum_{j \in J_n} \|E_j y_n\| \leq \theta_{m_{n_0}} \sum_{n < n_0} \# \text{supp } y_n \leq 1/2^{n_0}.$$

Notice that

$$\theta_m \sum_{j \in J_{n_0}} \|E_j y_{n_0}\| \leq \|y_{n_0}\| \leq 1$$

For $(E_j)_{j \in J_n, n \in L, n > n_t}$ choose (F_j) as in the Case 1. Thus we prove the Lemma with constant $C' = 3 \cdot 2^6 C$. $\square$

Proof of the Theorem 3.1. Take any block subspaces Y, Z of X and pick by Theorem 2.8 $(y_j) \subset Y$, $(z_j) \subset Z$ so that each $y_j = (y_1^j + \dots + y_{N_j}^j) / \|y_1^j + \dots + y_{N_j}^j\|$, $z_j = (z_1^j + \dots + z_{N_j}^j) / \|z_1^j + \dots + z_{N_j}^j\|$, where (y_n^j) , (z_n^j) are as in Lemmas 3.2, 3.3 and N_j are so big that $\|y_1^j + \dots + y_{N_j}^j\|, \|z_1^j + \dots + z_{N_j}^j\| \geq 2^{j+7}$ (for example $N_j \theta_{N_j} \geq 2^{j+7}$). Take any scalars $(a_j) \subset [0, 1]$ with $\|\sum_j a_j y_j\| = 1$. By Lemma 2.4 we pick a norming functional ϕ with $\phi(\sum_j a_j y_j) \geq 1/6$ and a tree-analysis $(\phi_t)_{t \in T}$ comparable with (y_n) . We will pick norming ψ with $\psi(\sum_j a_j z_j) \geq 1/2^4 C'$.

For any j pick t_j so that ϕ_{t_j} covers y_j . For any j let $\phi_1^j < \dots < \phi_{k_j}^j$ be all the successors of ϕ_{t_j} whose supports are contained in $\text{supp } y_j$. Then by Lemma 3.3 pick for any j suitable $F_1^j < \dots < F_{l_j}^j$ with $l_j \leq k_j$ satisfying (3.2). Replace in the tree-analysis of ϕ each ϕ_i^j by ψ_i^j so that $\|F_i^j z_j\| = \psi_i^j(z_j)$ and $\text{supp } \psi_i^j \subset \text{supp } z_j$. Then for any j

$$\theta_{k_j} \sum_{i=1}^{k_j} \phi_i^j(y_j) \leq C' \theta_{k_j} \sum_{i=1}^{l_j} \psi_i^j(z_j) + 6/2^{j+7}$$

hence

$$1/6 \leq \phi(\sum_j a_j y_j) \leq C' \psi(\sum_j a_j z_j) + 1/2^4$$

which ends the proof, since ψ is from the norming set. $\square$

Remark 3.4. Notice that the proof provides in every block subspace of X some subsequentially minimal block subspace. Indeed any block sequence of arbitrary long averages of vectors of the form $(y_1 + \dots + y_N) / \|y_1 + \dots + y_N\|$, where $y_1, \dots, y_N$ are as in Lemma 3.2 for arbitrary big N and $m_1, \dots, m_N$, spans a subsequentially minimal subspace.

Moreover by Lemma 3.2 sequences of long ℓ_p -averages generates equivalent spreading models. In case $p = 1$ the proof can be easily adapted to show that the spreading models of sequences of long ℓ_1 -averages are equivalent to the unit vector basis of the considered space X .

Now we consider special case: $\inf_n c_n = c > 0$. An example of such space is Tzafriri space $T[(\mathcal{A}_n, c/\sqrt{n})_n]$. Recall that a Banach space X with a basis belongs to Class 2 defined by T.Schlumprecht [22], if any block subspace Y of X contains normalized block sequences (x_n) , (y_n) such that the mapping carrying each x_n to y_n extends to a bounded strictly singular operator, i.e. operator whose no restriction to an infinite dimensional subspace is an isomorphism.

Theorem 3.5. *Let X be a p -space, $1 < p < \infty$, with $\inf_n c_n = c > 0$. Then*

- (1) *the dual space X^* is quasiminimal,*
- (2) *the spaces X and X^* do not contain a subspace of Class 2.*

Remark 3.6. By the above theorem the class of p -spaces with $\inf_n c_n > 0$ might provide an example of a Banach space containing neither a subspace of Class 2 nor a subspace of Class 1 (recall that a space with a basis is of Class 1 if any its normalized block sequence admits a subsequence equivalent to some subsequence of the basis, [22]), answering the question (Q4), [22].

In order to prove the theorem we need the following

Lemma 3.7. *Assume $\inf_n c_n = c > 0$. Take normalized block sequences (y_n) and (z_n) in X such that $\|z_n\|_\infty \leq c/16(\#\text{supp } y_n)^2$ for any n . Then (z_n) D -dominates (y_n) with some universal constant D .*

Proof of Lemma 3.7. We prove two Claims:

Claim 1. Fix $m \in \mathbb{N}$ and $1 > \delta > 0$. Then for any z with $\|z\| \geq 1/2$, $\|z\|_\infty < \delta/8m^2$ there are $F_1 < \dots < F_m$ such that

$$1 - \delta \leq \|F_i z\| / \|F_j z\| \leq 1 + \delta, \quad i, j = 1, \dots, m$$

Proof of Claim 1. Fix $\varepsilon > 0$. We prove induction on m that for any z with $\|z\|_\infty < \varepsilon$ and $m \leq \#\text{supp } z$ there are $F_1 < \dots < F_m$ such that $\sup_i \|F_i z\| \leq \inf_i \|F_i z\| + 2m\varepsilon$ and $F_1 \cup \dots \cup F_m = \text{supp } z$.

For $m = 2$ the claim easily follows. Assume the claim holds for $m - 1$. Take z with $\|z\|_\infty < \varepsilon$ and $\#\text{supp } z \geq m$. We can assume that $\text{supp } z = \{1, \dots, J\}$ for some $J \geq m$ to simplify the notation. For any $m - 1 \leq j \leq J$ let z_j be the restriction of z to the interval $\{1, \dots, j\}$. By the inductive assumption for $m - 1$ pick suitable $F_1^j < \dots < F_{m-1}^j$ for vectors z_j , $m - 1 \leq j \leq J$.

Notice that $\sup_i \|F_i^j z_j\| \leq \sup_i \|F_i^{j-1} z_{j-1}\| + (2m - 1)\varepsilon$. Indeed, if not, then by the choice of F_i^j 's we have $\|F_i^j z_j\| > \|F_k^{j-1} z_{j-1}\| + \varepsilon$ for any $1 \leq i, k \leq m - 1$, hence $\max F_i^j > \max F_i^{j-1} + 1$ for any $i \leq m - 1$, which by assumption on $\|z\|_\infty$ contradicts $\max \text{supp } z_j = \max \text{supp } z_{j-1} + 1$.

Notice also that $\|z - z_j\| + \varepsilon \geq \|z - z_{j-1}\|$ for any $m - 1 \leq j \leq J$.

Now if $\|z - z_{m-1}\| < \varepsilon$, then $\{1\}, \dots, \{m - 1\}, \text{supp}(z - z_{m-1})$ satisfy the claim.

Otherwise consider function $\xi(j) = \|z - z_j\| - \sup_i \|F_i^j z_j\|$ for $m - 1 \leq j \leq J$. We have that

$$\xi(m - 1) = \|z - z_{m-1}\| - \sup_i \|F_i^{m-1} z_{m-1}\| \geq \|z - z_{m-1}\| - \|z\|_\infty > 0$$

and $\xi(J) < 0$ and by previous observations $\xi(j) + 2m\varepsilon \geq \xi(j - 1)$ for any $m - 1 < j \leq J$. Take the minimal $J \geq j_0 > m - 1$ with $\xi(j_0) \leq 0$, then $0 < \xi(j_0 - 1) \leq \xi(j_0) + 2m\varepsilon$, hence

$$\sup_i \|F_i^{j_0} z_{j_0}\| - 2m\varepsilon \leq \|z - z_{j_0}\| \leq \sup_i \|F_i^{j_0} z_{j_0}\|$$

thus if we put $F_i = F_i^{j_0}$ for any $1 \leq i < m$ and $F_m = \text{supp}(z - z_{j_0})$, then we have by the inductive assumption and choice of j_0 that $\sup_{1 \leq i \leq m} \|F_i z\| \leq \inf_{1 \leq i \leq m} \|F_i z\| + 2m\varepsilon$.

Take $(F_i)_1^m$ as above. Take z as in Claim, then also $\#\text{supp } z \geq m$. Notice that $\sup \|F_i z\| \geq 1/2m$. Hence $\inf \|F_i z\| / \sup \|F_i z\| \geq 1 - 4m^2\varepsilon$ thus also $\sup \|F_i z\| / \inf \|F_i z\| \leq 1/(1 - 4m^2\varepsilon)$ which ends the proof. $\square$

Claim 2. Assume $\inf_n c_n = c > 0$. Take any normalized $y, z \in X$, assume that $\|z\|_\infty \leq c/16(\#\text{supp } y)^2$. Then for $m \in \mathbb{N}$, any $E_1 < \dots < E_m$ there are $F_1 < \dots < F_m$ with

$$\|E_1 y\| + \dots + \|E_m y\| \leq 9c^{-2}(\|F_1 z\| + \dots + \|F_m z\|)$$

Proof of Claim 2. Notice that by definition of the norm we have $\|E_1 y\| + \dots + \|E_m y\| \leq m^{1/q}/c$. Of course we can assume that $m \leq \#\text{supp } y$. Now we pick $F_1 < \dots < F_m$ by the Claim 1 for $\delta = 1/2$. Then by Lemma 2.10 we have

$$c\|z\|/6 \leq \left(\sum_i \|F_i z\|^p\right)^{1/p} \leq m^{1/p} 3 \inf_i \|F_i z\|/2$$

therefore $\|F_i z\| \geq c/9m^{1/p}$ for any i , hence

$$\sum_i \|F_i z\| \geq m^{1/q} c/9$$

which end the proof of the Claim. $\square$

This Claim shows in fact that Lemma 3.3 holds in X without error for much larger class of vectors provided the special assumption on (c_n) holds. Hence coming back to the proof of Lemma 3.7 we proceed as in the proof of Theorem 3.1, the situation is now simpler than in the general case, enabling us to pass to the dual space. Take any $a_1, \dots, a_N \in [0, 1]$. Take any functional ϕ from the norming set with a tree-analysis $(\phi_t)_{t \in T}$, assume by Lemma 2.4 that $(\phi_t)_{t \in T}$ is comparable with $y_1, \dots, y_N$. We will show that there is a functional ψ from the norming set such that $\phi(\sum_n a_n y_n) \leq 9c^{-4} \psi(\sum_n a_n z_n)$. For any n take t_n such that ϕ_{t_n} covers y_n and let $(\phi_i)_{i \in J_n}$ be immediate successors of ϕ_{t_n} with supports contained in $\text{supp } y_n$. Using Claim 2 find $(\psi_i)_{i \in J_n}$ with

$$\sum_{i \in J_n} \phi_i(y_n) \leq 9c^{-2} \sum_{i \in J_n} \psi_i(z_n)$$

and replace in the tree-analysis of ϕ all $(\phi_i)_{i \in J_n}$ with $(\psi_i)_{i \in J_n}$ obtaining in such a way a norming functional ψ with the desired property. $\square$

Proof of Theorem 3.5. We proceed to the proof of the first part. We need the following claims:

Claim 3. *For any $\delta, \alpha > 0$ there is some $\varepsilon > 0$ such that for any ϕ with $\|\phi\|_\infty < \varepsilon$ and any $x \in X$ with $\|x\| \leq 1$ and $\phi(x) > \alpha$ there is some $I \subset \mathbb{N}$ so that $|\phi(x_I)| \geq |\phi(x)|/2$ and $\|x_I\|_\infty < \delta$.*

Proof of Claim 3. Fix $\delta, \alpha > 0$. Take ϕ with $\|\phi\|_\infty < \varepsilon$ for some $\varepsilon > 0$ and x with $\|x\| \leq 1$ and put $J = \{i \in \text{supp } x : |x(i)| > \delta\}$. Then $c(\#J)^{1/p} \delta \leq \|x\| \leq 1$ from the definition of the norm in X , hence

$$|\phi(x_J)| = \sum_{j \in J} |\phi(j)x(j)| < \varepsilon \#J \leq \varepsilon / (c\delta)^p$$

Thus $\varepsilon = (c\delta)^p \alpha / 2$ and $I = \mathbb{N} \setminus J$ satisfy the desired property. $\square$

Claim 4. *Take normalized block sequences $(\phi_n), (\psi_n)$ in the norming set of X such that $\|\psi_n\|_\infty < \varepsilon_n$, where each ε_n , $n \in \mathbb{N}$, is chosen by Claim 3 for $\delta = c/4^{n+3}(\#\text{supp } \phi_n)^2$ and $\alpha = 1/4^n$.*

Then (ϕ_n) 4D-dominates (ψ_n) , with D as in Lemma 3.7.

Proof of Claim 4. Take $a_1, \dots, a_N$ and consider $\phi = a_1\phi_1 + \dots + a_N\phi_N$, $\psi = a_1\psi_1 + \dots + a_N\psi_N$. Let $\|\psi\| = 1$ and take $z \in X$ with $\|z\| = 1$ and $\psi(z) = 1$. We can assume, by unconditionality of the unit vector basis, that all coefficients of ϕ and z are non-negative, as well as $a_1, \dots, a_N$.

Let $J = \{n : \psi_n(z) \leq 1/4^n\}$. Then $\sum_{n \in J} a_n \psi_n(z) \leq 1/2$.

By the choice of (ε_n) we can take for any $n \notin J$ some vector w_n - a restriction of $z_{\text{supp } \psi_n}$ - such that $\|w_n\|_\infty \leq c/4^{n+3}(\#\text{supp } \phi_n)^2$ and $\|w_n\| \geq \psi_n(w_n) \geq \psi_n(z)/2 \geq 1/4^{n+1}$.

Take $v_n = w_n/\|w_n\|$, $n \notin J$, and notice that $\|v_n\|_\infty \leq c/16(\#\text{supp } \phi_n)^2$. Now pick normalized block sequence (y_n) with $\phi_n(y_n) = 1$, $\text{supp } y_n \subset \text{supp } \phi_n$. By Lemma 3.7 we have that (v_n) D-dominates (y_n) , hence the vector $y = \sum_{n \notin J} \|w_n\| y_n$ satisfies $\|y\| \leq D \|\sum_{n \notin J} w_n\| \leq D \|z\| = D$. Therefore

$$D \|\phi\| \geq \phi(y) = \sum_{n \notin J} a_n \|w_n\| \geq \sum_{n \notin J} a_n \psi_n(w_n) \geq \sum_{n \notin J} a_n \psi_n(z)/2 \geq 1/4$$

$\square$

Now take any block subspace $Y, Z \subset X^*$. Pick inductively infinite normalized block sequences $(\phi_n) \subset Y$, $(\psi_n) \subset Z$ such that

- (1) $\|\psi_n\|_\infty < \varepsilon_n$, where ε_n is chosen by Claim 3 for $\delta = c/4^{n+3}(\#\text{supp } \phi_n)^2$ and $\alpha = 1/4^n$, for any n ,
- (2) $\|\phi_{n+1}\|_\infty < \varepsilon'_n$, where ε'_n is chosen by Claim 3 for $\delta = c/4^{n+3}(\#\text{supp } \psi_n)^2$ and $\alpha = 1/4^n$, for any n .

Then by Claim 4, (ϕ_n) $4D$ -dominates (ψ_n) and (ψ_n) $4D$ -dominates (ϕ_{n+1}) . Now split $\mathbb{N}$ into intervals $I_1 < I_2 < \dots$ such that $\|\sum_{n \in I_j} \phi_n\| \geq 2^j$ and $\|\sum_{n \in I_j} \psi_n\| \geq 2^j$ (this is possible, since X^* does not contain c_0). The following Lemma ends the proof of the first part of the Theorem:

Lemma 3.8. *Let X be a Banach space. Take normalized 1-unconditional basic sequences $(u_n), (v_n) \subset X$ so that (u_n) dominates (v_n) , (v_{n+1}) dominates (u_n) and sequence $I_1 < I_2 < \dots$ of intervals of $\mathbb{N}$ such that $\|\sum_{n \in I_j} u_n\| \geq 2^j$, $\|\sum_{n \in I_j} v_n\| \geq 2^j$. Let $y_j = \sum_{n \in I_j} u_n / \|\sum_{n \in I_j} u_n\|$, $z_j = \sum_{n \in I_j} v_n / \|\sum_{n \in I_j} v_n\|$ for any $j \in \mathbb{N}$. Then (y_j) and (z_j) are equivalent.*

Proof of Lemma. Let $D \geq 1$ be the domination constant. Notice first that

$$\left\| \sum_{n \in I_j} v_n \right\| \leq D \left\| \sum_{n \in I_j} u_n \right\|$$

and

$$\left\| \sum_{n \in I_j} u_n \right\| \leq D \left\| \sum_{n \in I_j} v_{n+1} \right\| \leq D \left\| \sum_{n \in I_j} v_n \right\| + D \|v_{\max I_j + 1}\| \leq 2D \left\| \sum_{n \in I_j} v_n \right\|$$

Hence, by (u_n) dominating (v_n) , for any scalars (a_j) we have

$$\left\| \sum_j a_j z_j \right\| \leq 2D^2 \left\| \sum_j a_j y_j \right\|$$

To see the other domination take (a_j) with $\|\sum_j a_j z_j\| = 1$. Then, by (v_{n+1}) dominating (u_n) , we have

$$\begin{aligned} \left\| \sum_j a_j y_j \right\| &\leq \sum_j (|a_j| / \left\| \sum_{n \in I_j} u_n \right\|) + \left\| \sum_j a_j \left(\sum_{n \in I_j \setminus \{\max I_j\}} u_n \right) / \left\| \sum_{n \in I_j} u_n \right\| \right\| \\ &\leq 1 + 2D^2 \left\| \sum_j a_j z_j \right\| \leq 3D^2 \end{aligned}$$

□

In order to show the second part of the Theorem in case of space X we prove the following

Claim 5. *Let $\inf_n c_n = c > 0$. Take a normalized block sequences $(u_n), (v_n) \subset X$ with $\|u_n\|_\infty \rightarrow 0$ and $\|v_n\|_\infty \rightarrow 0$. Then if the mapping $v_n \rightarrow u_n$ extends to a bounded linear operator T then there is an infinite dimensional subspace Y of $[u_n]$ with $T|_Y$ an isomorphism.*

Proof of Claim. Take $(u_n), (v_n)$ as above. Assume that T defined as above is strictly singular. Pick inductively subsequence (u_{i_n}) and (v_{j_n}) such that $\|v_{j_n}\|_\infty < c/16(\#\text{supp } v_{i_n})$ and $\|u_{i_{n+1}}\|_\infty < c/16(\#\text{supp } v_{j_n})$. Then by Lemma 3.7 (v_{j_n}) dominates (v_{i_n}) , hence also (u_{i_n}) , and $(u_{i_{n+1}})$ dominates (v_{j_n}) . Denote the operator carrying each v_{i_n} to u_{i_n} by T' . As a composition of a strictly singular operator and a bounded right-hand shift operator T' is also strictly singular.

On the other hand we can consider averages $y_j = \sum_{n \in I_j} u_{i_n} / \|\sum_{n \in I_j} u_{i_n}\|$, $z_j = \sum_{n \in I_j} v_{i_n} / \|\sum_{n \in I_j} v_{i_n}\|$ for some sufficiently long (I_j) . Then by Lemma 3.8 $(y_j), (z_j)$ are equivalent and moreover

$$1/D' \leq \left\| \sum_{n \in I_j} u_{i_n} \right\| / \left\| \sum_{n \in I_j} v_{i_n} \right\| \leq D'$$

for some $D' \geq 1$. Hence T' restricted to $[z_j]$ is an isomorphism, which gives contradiction. $\square$

The case of the dual space X^* follows analogously, by using Claim 4. $\square$

4. QUASIMINIMALITY IN MIXED TSIRELSON SPACES $T[(\mathcal{S}_n, \theta_n)_n]$

Notice ([2]) that any mixed Tsirelson space $T[(\mathcal{S}_{k_n}, \theta_n)_{n \in \mathbb{N}}]$, with $\theta \rightarrow 0$, is isometric to $T[(\mathcal{S}_n, \widehat{\theta}_n)_{n \in \mathbb{N}}]$, where

$$\widehat{\theta}_n = \sup \left\{ \prod_{i=1}^l \theta_{n_i} : \sum_{i=1}^l n_i \geq n \right\}, \quad n \in \mathbb{N}$$

Notice that in such case the sequence $(\widehat{\theta}_n)$ satisfies $\widehat{\theta}_n \searrow 0$ and $\widehat{\theta}_{n+m} \geq \widehat{\theta}_n \widehat{\theta}_m$ for any $n, m \in \mathbb{N}$. We will assume in the paper that we work in such a setting.

Definition 4.1. [2] *A mixed Tsirelson space $T[(\mathcal{S}_n, \theta_n)_{n \in \mathbb{N}}]$ is called regular if $\theta_n \searrow 0$ and $\theta_{n+m} \geq \theta_n \theta_m$ for any $n, m \in \mathbb{N}$.*

Notation. Take a regular mixed Tsirelson space $T[(\mathcal{S}_n, \theta_n)_{n=1}^\infty]$. Then there exists $\theta = \lim_n \theta_n^{1/n} = \sup_n \theta_n^{1/n} \in (0, 1]$. We will use the following notation: $\theta_n = c_n \theta^n$, with $c_n \in (0, 1]$, $n \in \mathbb{N}$. If $c_n = 1$ for some $n \in \mathbb{N}$ then $T[(\mathcal{S}_n, \theta_n)]$ is isomorphic to Tsirelson space $T[\mathcal{S}_1, \theta]$.

4.1. Two lemmas for the Schreier families. We prove first technical lemmas needed in the sequel.

Lemma 4.2. *Let $k, m \in \mathbb{N}$ and $l \in \mathbb{N}$ such that $km < 2^l$. Then*

$$(\mathcal{S}_n[\mathcal{A}_k])[\mathcal{A}_m] \subset \mathcal{A}_l[\mathcal{S}_n] \text{ for any } n \in \mathbb{N}.$$

Proof. We prove the result by induction on n . For $n = 1$ we have

$$[\mathcal{S}_1[\mathcal{A}_k]][\mathcal{A}_m] = \mathcal{S}_1[\mathcal{A}_k[\mathcal{A}_m]] = \mathcal{S}_1[\mathcal{A}_{km}]$$

Let $j \leq m_1 < \dots < m_j$ and

$$m_i \leq \lambda_{(i-1)km+1} < \dots < \lambda_{(i-1)km+km-1} < \lambda_{kmi} < m_{i+1}$$

for every $i = 1, \dots, j$. We set $G = \{\lambda_1, \dots, \lambda_{kmj}\}$. From the assumptions we have that $j \leq G$.

For $i = 1, \dots, l$ consider the sets

$$F_i = \{2^{i-1}j, 2^{i-1}j + 1, \dots, 2^i j - 1\}$$

Each of the sets F_i , $i \leq l$ belongs to the family $\mathcal{S}_1$ since $\#F_i \leq \min F_i$. We may write $G \subset \bigcup_{i=1}^l G_i$ where $G_i = \{\lambda_{r-j+1} : r \in F_i\}$ for $i \leq l$. Then each G_i is a spreading of F_i and hence belongs to the Schreier family $\mathcal{S}_1$. Thus $G \in \mathcal{A}_l[\mathcal{S}_1]$.

Assume that the result holds for n , i.e. $[\mathcal{S}_n[\mathcal{A}_k]][\mathcal{A}_m] \subset \mathcal{A}_l[\mathcal{S}_n]$. Then we have that

$$[\mathcal{S}_{n+1}[\mathcal{A}_k]][\mathcal{A}_m] = \mathcal{S}_1[\mathcal{S}_n[\mathcal{A}_k]][\mathcal{A}_m] \subset \mathcal{S}_1[\mathcal{A}_l[\mathcal{S}_n]] \subset \mathcal{A}_l[\mathcal{S}_{n+1}]$$

The last inclusion follows immediately from the associativity of the operation $M[N]$ and the fact that $\mathcal{A}_l[\mathcal{S}_n] = (\mathcal{S}_n)^l$. $\square$

Lemma 4.3. *Let $F \in \mathcal{S}_m$ and $(F_i)_{i=1}^d$ be a partition of F into successive sets such that for every $i \leq d$, $F_i \in \mathcal{S}_{m_i} \setminus \mathcal{S}_{m_i-1}$ for some $m_i \leq m$. For every $i \leq d$ let $G_i \in \mathcal{S}_{m_i-1}$ (in case $m_i = 0$ let $G_i \in \mathcal{S}_0$) with $G_1 < \dots < G_d$. Then we have the following:*

- (1) *if $3 \max F_i < \min G_{i+1}$ for every $i = 1, \dots, d-1$, then $\bigcup_{i=2}^d G_i \in \mathcal{S}_m$ and $\bigcup_{i=1}^d G_i \in \mathcal{A}_2[\mathcal{S}_m]$.*
- (2) *if $2 \max F_i < \min G_i$ for every $i = 1, \dots, d$, then $\bigcup_{i \leq d} G_i \in \mathcal{S}_m$.*

Proof. We prove the first part by induction on m . For $m = 1$ the result is clear since $G_i \in \mathcal{S}_0$ and also for $i \geq 2$, $G_i > \min F$.

Assume that the statement holds for m , and let $F \in \mathcal{S}_{m+1}$ and $(F_i)_{i=1}^d$ be a partition of F into successive subsets.

We have also that $F = \bigcup_{i=1}^k A_i$ where $A_i \in \mathcal{S}_m$ and $k \leq A_1 < A_2 < \dots < A_k$. Let $C_i = \{j \in \{2, 3, \dots, d\} : F_j \subset A_i\}$ for $i = 1, \dots, k$. We have that $F_j \in \mathcal{S}_m$ for every j with $F_j \subset A_i$ for some i . From inductive hypothesis we get that

$$U_i = \bigcup_{j \in C_i} G_j \in \mathcal{A}_2[\mathcal{S}_m] \text{ for every } i \leq k.$$

Let $U_i = U_{i,1} \cup U_{i,2}$ with $U_{i,1} < U_{i,2}$ in $\mathcal{S}_m$. Let us observe that we have $3k \leq U_{i,1}$ for every $i = 1, \dots, k$.

Consider also $B = \{j \leq d : F_j \text{ is not contained in any } \mathcal{A}_i\}$. It follows that $\#B \leq k$ and therefore $\mathcal{L} = \{U_{i,1}, U_{i,2} : i = 1, \dots, k\} \cup \{G_j : j \in B\}$ contains at most $3k$ elements of $\mathcal{S}_m$ with $3k \leq \text{minsupp } G_2 \leq \text{minsupp } \mathcal{L}$ and therefore $\bigcup \mathcal{L} \in \mathcal{S}_{m+1}$.

Since $\mathcal{L}$ contains all G_j except G_1 we get that

$$\bigcup \{G_j : j \leq d\} \in \mathcal{A}_2[\mathcal{S}_{m+1}].$$

The proof of the second part follows similarly, in a shorter way, since we don't need the partition of sets U_i into elements of $\mathcal{S}_m$ and treating separately the set G_1 . $\square$

4.2. The quasiminimality. We assume that in the whole section we work with a regular mixed Tsirelson space $X = T[(\mathcal{S}_n, \theta_n)_{n \in \mathbb{N}}]$.

The arguments to prove the quasiminimality of the space X will follow the arguments we use for the p -spaces, showing the existence of special averages and forming from them equivalent sequences.

We will need averages of a certain type, playing the role of ℓ_p -averages in p -spaces. The major tool providing these averages is Lemma 4.11 [2] stated in general case, however with a proof in [2] requiring an additional assumption - that the sequence $(c_n)_n$ is decreasing. Thus we need this assumption in the proof of Theorem 4.10, and we use it only when applying Lemma 4.11 [2]. Therefore any mixed Tsirelson space $T[(\mathcal{S}_n, \theta_n)_n]$ in which Lemma 4.11 holds, is quasiminimal.

Let us mention that as observed by the first named author and D. Leung (private communication) the proof of the main result of [2] on distortion of spaces $T[(\mathcal{S}_n, \theta_n)_n]$ with $c_n \rightarrow 0$ can be adapted to work in all cases just following the proof of the main result from the preprint version of [2] in arxiv.org.

In order to have an analogue to Lemma 2.4 we shall use the auxiliary space $X_k = T[(\mathcal{S}_n[\mathcal{A}_k], \theta_n)_n]$, $k \in \mathbb{N}$. It follows readily from the definition that $\|\cdot\|_X \leq \|\cdot\|_{X_k}$. In the next lemma we prove that the spaces are isomorphic.

Lemma 4.4. *The spaces X and X_k are $(k+1)$ -isomorphic.*

In particular for every functional f in the norming set of X_k there exists $(k+1)$ functionals $\phi_1 < \dots < \phi_{k+1}$ in the norming set of X such that $f = \phi_1 + \dots + \phi_{k+1}$ and the restriction of the tree-analysis of f to the support of ϕ_i gives a tree-analysis of ϕ_i , $i = 1, \dots, k+1$.

Proof. We prove it by induction on the height of the tree-analysis. If $f = \theta_n \sum_{i \in A} e_i$, with $A \in \mathcal{S}_n[\mathcal{A}_k]$, then by Lemma 4.2 $A \in \mathcal{A}_{k+1}[\mathcal{S}_n]$ hence $A = A_1 \cup \dots \cup A_{k+1}$ for some A_t 's from $\mathcal{S}_n$, hence $\phi_t = \theta_n \sum_{i \in A_t} e_i$, $t = 1, \dots, k+1$, give the desired partition.

Now take $f = \theta_n \sum_{i \in A} f_i$ for some $A \in \mathcal{S}_n[\mathcal{A}_k]$ and some norming functionals (f_i) of X_k . By inductive assumption

$$f = \theta_n \sum_{i \in A} (\phi_{i,1} + \cdots + \phi_{i,k+1})$$

for some norming functionals $(\phi_{i,j})_{i \in A, j=1, \dots, k+1}$ of X .

Since $(f_i)_{i \in A}$ is $\mathcal{S}_n[\mathcal{A}_k]$ -admissible and $k(k+1) < 2^{k+1}$, Lemma 4.2 yields $(\phi_{i,j})_{i \in A, j=1, \dots, k+1}$ is $\mathcal{A}_{k+1}[\mathcal{S}_n]$ -admissible. Therefore there are some $G_1, \dots, G_{k+1}$ such that $(\phi_{i,j})_{(i,j) \in G_t}$ is $\mathcal{S}_n$ -admissible and setting $\phi_t = \theta_n \sum_{(i,j) \in G_t} \phi_{i,j}$ for $t = 1, \dots, k+1$ we get $(k+1)$ functionals $\phi_1 < \cdots < \phi_{k+1}$ in the norming set of X with $\phi_1 + \cdots + \phi_{k+1} = f$.

It follows that $|f(u)| \leq (k+1) \max\{|\phi_1(u)|, \dots, |\phi_{k+1}(u)|\}$ for any $u \in X$. The restriction of the tree-analysis of f to the support of each ϕ_t gives the tree-analysis of ϕ_t in X with the desired property. $\square$

We prove now an analogue to Lemma 2.4.

Lemma 4.5. *Take a functional ϕ from the norming set of X and any finite block sequence (v_n) . Set $v = \sum_n v_n$. Then there is a functional ϕ' with the tree-analysis $(\phi'_t)_{t \in T'}$ comparable with (v_n) and $4\phi'(v) \geq \phi(v)$.*

Proof. For the proof we consider the auxiliary space X_3 . Take ϕ in the norming set of X with a tree-analysis $(\phi_t)_{t \in T}$. For any $t \in T$ set $I_t = \{n : v_n \text{ is covered by } \phi_t\}$ and set for every $n \in I_t$, $J_{t,n} = \{r \in \text{succ } t : \text{range } \phi_r \cap \text{range } v_n \neq \emptyset\}$.

Observe that every $r \in \text{succ } t$ belong to at most two sets $J_{t,n}$, say J_{t,n_1}, J_{t,n_2} . So we can split each $\text{supp } \phi_r$ into three norming functionals of X with a tree-analysis comparable to (v_n) : $(\phi_r)|_{\text{supp } v_{n_1}}$, $(\phi_r)|_{\text{supp } v_{n_2}}$ and the remaining part.

Applying the above argument for all $t \in T$ with $I_t \neq \emptyset$ and $r \in \text{succ } t$, starting from the root and moving “downward” the tree, we produce a tree-analysis comparable with (v_n) of some functional f in the norming set of X_3 .

By Lemma 4.4 the functional f of X_3 can be represented as the sum of four norming functionals $\phi_1 < \phi_2 < \phi_3 < \phi_4$ of X such that restriction of the tree-analysis of f to the support of each ϕ_l gives the tree-analysis of ϕ_l . Therefore each ϕ_l has a tree-analysis comparable with (v_n) . To end the proof pick ϕ' from $\{\phi_1, \dots, \phi_4\}$ satisfying $\phi'(v) \geq \phi(v)/4$. $\square$

We recall now the notion of averages given in [2], Definition 4.5, with simplified notation. Averages of this type, called special convex combinations, were also used in studies of mixed Tsirelson spaces in particular in [4, 5, 14, 13].

Definition 4.6. [2] *Fix $M \in \mathbb{N}$, $\varepsilon > 0$ and take a block sequence $(x_n) \subset X$. We say that $x \in X$ is an (M, ε) average of (x_n) given by an averaging tree $T = (x^{i,j})_{j=0, i=1}^{M, N_j}$ on X if*

- (1) $1 = N_M < N_{M-1} < \cdots < N_0$,
- (2) $x = x^{1,M}$ is the root of T , the sequence $(x^{i,0})_{i=1}^{N_0}$ of maximal nodes of T form a subsequence of (x_n) ,
- (3) $\text{succ } x^{i,j} = (x^{s,j-1})_{s \in I_{i,j}}$ for some non-empty interval $I_{i,j} \subset \{1, \dots, N_j\}$ with $(x^{s,j-1})_{s \in I_{i,j}}$ a block sequence $\mathcal{S}_1$ -admissible with respect to (x_n) , for any $j = 0, \dots, M$, $i = 1, \dots, N_j$,
- (4) $x^{i,j} = \frac{1}{k_{i,j}} \sum_{s \in I_{i,j}} x^{s,j-1}$, where $k_{i,j} = \#I_{i,j}$, for any $j = 0, \dots, M$, $i = 1, \dots, N_j$,
- (5) $k_{1,j} > 6 \cdot 2^{2+j} \theta^{-1} \varepsilon^{-1}$, $k_{i,j} > 6 \cdot 2^{1+i+j} \theta^{-1} \varepsilon^{-1} \max \text{supp } x^{i-1,j}$ for any $i = 2, \dots, N_j$, $j = 0, \dots, M$.

Remark 4.7. Recall that by Proposition 4.7, [2], for any normalized block sequence (x_n) , $M \in \mathbb{N}$, $\varepsilon > 0$ there is a (M, ε) -average of (x_n) .

Lemma 4.8. *For any block subspace Y of X , any $\delta > 0$, $\varepsilon > 0$ and any $M \in \mathbb{N}$ there is a normalized block sequence $(x_i) \subset Y$ and a (M, ε) -average x of (x_i) given by an averaging tree $T = (x^{i,j})_{j=0,i=1}^{M,N_j}$ such that for any $j = 1, \dots, M$ we have*

$$\|x^{i,j}\| \geq (1 - \delta)^j \theta^j \quad i = 1, \dots, N_j$$

Proof. We use the standard argument as in Lemma 2.8 [4], Lemma 4.12 [2]. Assume the contrary and pick any normalized block sequence (x_n) in Y and let $(y_{n,1})$ be a block sequence of (M, ε) -averages of (x_n) . Let any $y_{n,1}$ be given by a tree $(x_{n,1}^{i,j})_{i,j}$. For any n there is some $1 \leq j_{n,1} \leq M$ and $i_{n,1}$ so that $\|x_{n,1}^{i_{n,1}, j_{n,1}}\| \leq (1 - \delta)^{j_{n,1}} \theta^{j_{n,1}}$. Hence there is an infinite $N_1 \subset \mathbb{N}$ and some $1 \leq J_1 \leq M$ so that $j_{n,1} = J_1$ for any $n \in N_1$.

Let $z_n = x_{n,1}^{i_{n,1}, J_1} / \|x_{n,1}^{i_{n,1}, J_1}\|$ for any $n \in N_1$. Let $(y_{n,2})$ be a block sequence of (M, ε) -averages of $(z_n)_{n \in N_1}$, each $y_{n,2}$ given by an averaging tree $(x_{n,2}^{i,j})_{i,j}$. Find some infinite $N_2 \subset N_1$, some sequence of integers $(i_{n,2})$ and some $1 \leq J_2 \leq M$ so that $\|x_{n,2}^{i_{n,2}, J_2}\| \leq (1 - \delta)^{J_2} \theta^{J_2}$ for any $n \in N_2$. Notice that by definition of averages each $x_{n,2}^{i_{n,2}, J_2}$ is a convex combination of some $\mathcal{S}_{J_2}$ -admissible z_n 's, and recall that any $x_{n,1}^{i,j}$ is a convex combination of some $\mathcal{S}_{J_1}$ -admissible x_n 's, so $x_{n,2}^{i_{n,2}, J_2} = \sum_{k \in F_{n,2}} a_k x_k$, for $(x_k)_{k \in F_{n,2}}$ $\mathcal{S}_{J_1+J_2}$ -admissible and $\sum_{k \in F_{n,2}} a_k \geq (1 - \delta)^{-J_1} \theta^{-J_1}$. Normalize $(x_{n,2}^{i_{n,2}, J_2})$ and continue in this manner.

After m steps we end up with a vector $x = x_{1,m}^{i_{1,m}, J_m}$ (first of some sequence) so that $\|x\| = \|x_{1,m}^{i_{1,m}, J_m}\| \leq (1 - \delta)^{J_m} \theta^{J_m}$. On the other hand, $x = \sum_{k \in F_{1,m}} a_k x_k$, for some $(x_k)_{k \in F_{1,m}}$ $\mathcal{S}_{J_1+\dots+J_m}$ -admissible and

$$\sum_{k \in F_{1,m}} a_k \geq (1 - \delta)^{-(J_1+\dots+J_{m-1})} \theta^{-(J_1+\dots+J_{m-1})}$$

Hence we get

$$(1 - \delta)^{J_m} \theta^{J_m} \geq \theta_{J_1+\dots+J_m} (1 - \delta)^{-(J_1+\dots+J_{m-1})} \theta^{-(J_1+\dots+J_{m-1})}$$

which yields $\theta_{J_1+\dots+J_m} \leq (1 - \delta)^{J_1+\dots+J_m} \theta^{J_1+\dots+J_m}$, which since $J_i \geq 1$, for large m contradicts the definition of θ . $\square$

Now we are ready to prove the existence of special (M, ε) -averages in all block subspaces of a mixed Tsirelson space. Recall that this fact is well-known and used in case of $\theta = 1$, e.g. [4, 5, 17, 14, 13].

Corollary 4.9. *Let $X = T[(\mathcal{S}_n, \theta_n)_n]$ be a regular mixed Tsirelson space with the sequence $(\frac{\theta_n}{\theta^n})_n$ decreasing. Then for any block subspace Y of X and any $M \in \mathbb{N}$ there is a vector $y \in Y$ with $\|y\| = 1$ such that for any $0 \leq j \leq M$*

$$4\theta_1^{-1} \theta^{-j-1} \geq \sup \left\{ \sum_i \|E_i y\| : (E_i) - \mathcal{S}_j \text{ admissible} \right\} \geq \theta_1 \theta^{1-j} / 4$$

Proof. Pick the (M, ε) -average x of (x_n) provided by Lemma 4.8, for δ with $(1 - \delta)^M \geq 1/2$ and $\varepsilon \leq \theta^{M-1} \theta_M$. It is enough to show that

$$2\theta^{M-1} \geq \|x\| \geq \theta_1 \theta^{M-1} / 2$$

$$2\theta^{M-j-1} \geq \sup \left\{ \sum_i \|E_i x\| : (E_i) - \mathcal{S}_j \text{ admissible} \right\} \geq \theta^{M-j} / 2, \quad j = 1, \dots, M$$

The lower estimate for $j = 1, \dots, M$ is provided by Lemma 4.8. Let x be given by a tree $(x^{i,j})_{i,j}$. Notice that for any $j = 1, \dots, M-1$ we have $x = \sum_{i=1}^{N_{M-j}} a_i^j x^{i, M-j}$

for some $(a_i^j) \subset [0, 1]$ with $\sum_{i=1}^{N_{M-j}} a_i^j = 1$ and $(x^{i, M-j})_{i=1}^{N_{M-j}}$ is $\mathcal{S}_j$ -admissible, which ends the proof. For $j = 0$ it is enough to use estimation for $j = 1$ and definition of the norm.

The upper estimate for any $j = 1, \dots, M$ follows from the proof of Lemma 4.11 (7), [2], in case $(\frac{\theta_n}{\theta^n})_n$ decreasing, for $N = 1$. The upper estimate for $j = 0$, i.e. the estimate of $\|x\|$ follows again from the proof Lemma 4.11 (6), [2], in case $(\frac{\theta_n}{\theta^n})_n$ decreasing, for $N = 1, J = M, i = 1$. $\square$

Thanks to more complex structure of families $(\mathcal{S}_n)_{n \in \mathbb{N}}$, which can "absorb" families $\mathcal{A}_n$ as it is reflected in Lemma 4.2, the construction of equivalent sequences in the spaces $T[(\mathcal{S}_n, \theta_n)_{n \in \mathbb{N}}]$ is simpler than the one in case of p -spaces. Namely in p -spaces case we produce equivalent sequences from averages of ℓ_p -averages. Now the equivalent sequence consist simply of "special averages" provided by Corollary 4.9.

Theorem 4.10. *Any regular mixed Tsirelson space $X = T[(\mathcal{S}_n, \theta_n)_{n \in \mathbb{N}}]$, with the sequence $(\frac{\theta_n}{\theta^n})_n$ decreasing, is quasiminimal.*

By Corollary 4.9 in order to prove the Theorem it is enough to show the following

Lemma 4.11. *Let $(y_n), (z_n) \subset X$ be normalized block sequences satisfying for some $(m_n) \subset \mathbb{N}$ the following:*

- (1) $2 \maxsupp y_n < z_n$ and $3 \maxsupp z_n < y_{n+1}$ for any $n \in \mathbb{N}$,
- (2) y_n and z_n satisfies the conclusion of Corollary 4.9 with $M = m_n$ for every $n \in \mathbb{N}$,
- (3) $\theta_{m_{n+1}} \maxsupp z_n < 2^{-n}$, $\theta_{m_{n+1}} \maxsupp y_n < 2^{-n}$ for any $n \in \mathbb{N}$.

Then $(y_n)_n$ and $(z_n)_n$ are equivalent.

Proof of Lemma 4.11. We show first that $(z_n)_n$ dominates $(y_n)_n$ with constant $C = (4 \cdot 2) \cdot (2 \cdot \frac{4^2}{\theta_1^2 \theta^3})$. Take any scalars $(a_n)_{n \in D}$. We shall use again the auxiliary space $X_2 = T[(\mathcal{S}_n[A_2], \theta_n)_n]$. Namely for every functional ϕ in the norming set of X with a tree-analysis $(\phi_t)_{t \in T}$ comparable with (y_n) we shall produce two functionals f_1, f_2 in the norming set of X_2 such that

$$(4.1) \quad \phi(\sum_{n \in D} a_n y_n) \leq (2 \cdot \frac{4^2}{\theta_1^2 \theta^3})(f_1 + f_2)(\sum_{n \in D} a_n z_n)$$

By Lemma 4.4 and Lemma 4.5 this will prove that $\|\sum_n a_n y_n\| \leq C \|\sum_n a_n z_n\|$.

For every $t \in T$ we set

$$D_t = \{n \in \mathbb{N} : \text{supp } \phi_t \cap \text{supp } y_n = \text{supp } \phi \cap \text{supp } y_n\}$$

and as usual $I_t = \{n \in D_t : \phi_t \text{ covers } y_n\}$.

We shall produce inductively (on level of $t \in T$) for any $t \in T$ two functionals $f_{t,1} < f_{t,2}$ in the norming set of X_2 with supports contained in $\bigcup \{\text{supp } z_n : n \in D_t\}$ such that (4.1) holds for ϕ_t, D_t and $f_{t,1}, f_{t,2}$.

Assume t is maximal, then $\phi_t = e_i^*$ for some n_0 . If $D_t = \emptyset$ let $f_{t,1} = f_{t,2} = 0$, otherwise $D_t = I_t = \{n\}$ and let $f_{t,1}$ be the norming functional of z_n in X , $f_{t,2} = 0$.

Assume that we have chosen suitable pairs of functionals for any functionals on the levels $l+1, l+2, \dots, \text{height}(T)$ and take ϕ_t on level l . Let $\phi_t = \theta_k \sum_{s \in \text{succ } t} \phi_s$. As usual for $n \in I_t$ we set $J_n = \{s \in \text{succ } t : \text{supp } \phi_s \subset \text{range } y_n\}$.

CASE 1. $k \leq m_n$ for every $n \in I_t$.

Then if $(\phi_s)_{s \in J_n}$ is $\mathcal{S}_{k_n}$ - and not $\mathcal{S}_{k_n-1}$ -admissible by condition 2. in choice of (y_n) we get

$$\sum_{s \in J_n} \phi_s(y_n) \leq 4\theta_1^{-1} \theta^{-k_n-1}$$

Hence by condition 2. in choice of (z_n) we can take an $\mathcal{S}_{k-1}$ -admissible set of functionals $(\psi_u)_{u \in V_n}$ from the norming set of X with $\text{supp } \psi_u \subset \text{supp } z_n$, $u \in V_n$, such that

$$\sum_{s \in J_n} \phi_s(y_n) \leq \frac{4^2}{\theta_1^2 \theta^3} \sum_{u \in V_n} \psi_u(z_n)$$

>From the inductive hypothesis for every $s \in \text{succ } t$ such that $D_s \neq \emptyset$ we have take two suitable functionals $f_{s,1} < f_{s,2}$ with $\text{supp } f_{s,1}, \text{supp } f_{s,2} \subset \{\text{supp } z_n : n \in D_s\}$. From Lemma 4.3(b) we get that the set

$$\{\text{supp } \psi_u : u \in V_n, n \in I_t\} \cup \{\text{supp } f_{s,1} : D_s \neq \emptyset\}$$

is $\mathcal{S}_k$ -admissible and therefore the sequence

$$\bigcup_{n \in I_t} \{\psi_u : u \in V_n\} \cup \bigcup_{s \in \text{succ } t : D_s \neq \emptyset} \{f_{s,1}, f_{s,2}\}$$

is $\mathcal{S}_k[\mathcal{A}_2]$ -admissible. It follows that

$$f_{t,1} = \theta_k \left(\sum_n \sum_{u \in V_n} \psi_u + \sum_{s : D_{\phi_s} \neq \emptyset} (f_{s,1} + f_{s,2}) \right)$$

is in the norming set of the set of the space X_2 and

$$\phi_t \left(\sum_{n \in D_t} y_n \right) \leq \frac{4^2}{\theta_1^2 \theta^3} f_{t,1} \left(\sum_{n \in D_t} a_n z_n \right).$$

In this case we set $f_{t,2} = 0$.

CASE 2. There exists $n \in I_t$ such that $m_n < k$.

Then let n_0 be the unique $n \in I_t$ such that $m_n \leq k < m_{n+1}$. For every $n \in D_t$ with $n < n_0$ we have $\theta_k \text{maxsupp}(y_n) \leq 2^{-n}$ so with error $\max |a_n| \sum_{n \in D_t, n < n_0} 2^{-n}$ we can erase this part.

As usual we compare $\phi_t(a_{n_0} y_{n_0})$ and $\phi_t(\sum_{I_t \ni n > n_0} a_n y_n)$. If the second term dominates the first one we proceed as in Case 1, but multiplying by 2 the estimation of $\phi_t(\sum_{n \in I_t} a_n y_n)$.

If the first term dominates the second one, take the functional

$$f_{t,2} = \theta_k \sum_{s \in \text{succ } t : D_s \neq \emptyset} (f_{s,1} + f_{s,2}).$$

Since the set $\{\phi_s : D_s \neq \emptyset\}$ is $\mathcal{S}_k$ -admissible, it is clear that $f_{t,2}$ is in the norming set of X_2 . Now take a functional $f_{t,1}$ that norms z_{n_0} in X . Of course we may assume that $\text{range } f_{t,1} \subset \text{range } z_{n_0}$. We also have that $f_{t,1} < f_{t,2}$ since we have delete all z_n for $n \in D_t$ and $n < n_0$. Hence we have produce two functionals $f_{t,1} < f_{t,2}$ in the norming set of X_2 satisfying the desired conditions.

We prove now that $(y_n)_n$ dominates $(z_n)_n$ in an analogous way.

Fix scalars $(a_n)_{n \in D}$. Take any norming functional ϕ and assume by Lemma 4.5 that ϕ has a tree-analysis $(\phi_t)_{t \in T}$ comparable with (z_n) .

For every $t \in T$ with $D_t = \{n : \text{supp } \phi_t \cap \text{supp } z_n = \text{supp } \phi \cap \text{supp } z_n\} \neq \emptyset$ we shall produce two functionals $f_{t,1} < f_{t,2}$ in the norming set of the space X_2 such that $\text{supp } f_{t,1}, \text{supp } f_{t,2} \subset \bigcup \{\text{supp } y_n : n \in D_t\}$ and

$$\phi_t \left(\sum_{n \in D_t} a_n z_n \right) \leq C(f_{t,1} + f_{t,2}) \left(\sum_{n \in D_t} a_n y_n \right)$$

If t is maximal we proceed as in the previous part. Assume we have done it for all $t \in T$ in levels $l+1, \dots, \text{height}(T)$. Let $\phi_t = \theta_k \sum_{s \in \text{succ } t} \phi_s$, for some $\{\phi_s : s \in \text{succ } t\} - \mathcal{S}_k$ -admissible.

We set

$$I_t = \{n \in D_t : \phi_t \text{ covers } z_n\} \text{ and } B = \{s \in \text{succ } t : D_s \neq \emptyset\}.$$

For every $n \in I_t$ we set $J_n = \{s \in \text{succ } t : \text{supp } \phi_s \subset \text{range } y_n\}$.

CASE 1. $k \leq m_n$ for every $n \in I_t$.

We have that $\text{succ } t = \bigcup_{n \in I_t} \{s : s \in J_n\} \cup \bigcup_{s \in B} \{s\}$. By condition 2. of choice of (y_n) and (z_n) for every $n \in I_t$ we choose $\{\psi_u : u \in V_n\}$ and $\mathcal{S}_{k_n-1}$ -admissible set, where $\mathcal{S}_{k_n}$ is the admissibility of the set $\{\phi_s : s \in J_n\}$, such that

$$\sum_{s \in J_n} \phi_s(z_n) \leq \frac{4^2}{\theta_1^2 \theta^3} \sum_{u \in V_n} \psi_u(y_n).$$

By Lemma 4.3 we get that the set

$$\bigcup_{n \in I_t} \{\psi_u : u \in V_n\} \cup \{f_{s,1} : s \in B\}$$

is $\mathcal{A}_2[\mathcal{S}_k]$ -admissible. Therefore we have partition the set

$$\begin{aligned} & \bigcup_{n \in I_t} \{\psi_u : u \in V_n\} \cup \{f_{s,1} : s \in B\} \\ &= (\{\psi_u : u \in V_1\} \cup \{f_{s,1} : s \in B_1\}) \cup (\{\psi_u : u \in V_2\} \cup \{f_{s,1} : s \in B_2\}) \end{aligned}$$

into two successive $\mathcal{S}_k$ -admissible sequences. Therefore the functionals

$$f_{t,i} = \theta_k \left(\sum_{u \in V_i} \psi_u + \sum_{s \in B_i} (f_{s,1} + f_{s,2}) \right), \quad i = 1, 2$$

are from the norming set of X_2 , satisfying the desired estimation. Indeed

$$\begin{aligned} \phi_t \left(\sum_{n \in D_t} a_n z_n \right) &= \theta_k \sum_{n \in I_t} \sum_{s \in J_n} \phi_s(a_n z_n) + \theta_k \sum_{s \in B} \phi_s \left(\sum_{n \in D_s} a_n z_n \right) \\ &\leq \theta_k \sum_{n \in I_t} \sum_{u \in V_n} C \psi_u(a_n y_n) + \theta_k \sum_{s \in B} C(f_{s,1} + f_{s,2}) \left(\sum_{n \in D_s} a_n y_n \right) \\ &= C(f_{t,1} + f_{t,2}) \left(\sum_{n \in D_t} a_n y_n \right). \end{aligned}$$

CASE 2. There exist $n \in I_t$ such that $m_n \leq k$.

Take $n_0 = \max\{n : m_n \leq k\}$. By condition 3. in the choice of the sequences $(y_n), (z_n)$ we can erase the $y_n, n < n_0, n \in D_t$ with error $\sum_{n \in D_t, n < n_0} |a_n| 2^{-n}$.

Now we compare $\phi_t(a_{n_0} z_{n_0})$ and $\theta_k \sum_{I_t \ni n > n_0} \sum_{s \in J_n} \phi_s(a_n z_n)$. Assume that the first term dominates the second. It follows

$$\phi_t(a_{n_0} z_{n_0}) + \theta_k \sum_{I_t \ni n > n_0} \sum_{s \in J_n} \phi_s(a_n z_n) \leq 2\phi_t(a_{n_0} z_{n_0}).$$

In this case we take a functional $f_{t,1}$ that norms y_{n_0} and we set

$$f_{t,2} = \theta_k \sum_{s \in B} (f_{s,1} + f_{s,2}).$$

It is clear that $f_{t,1} < f_{t,2}$ and by Lemma 4.3 we get that $\{f_{s,1} : s \in B\}$ is $\mathcal{S}_k$ -admissible hence $f_{t,2}$ is in the norming set of X_2 . Therefore

$$\begin{aligned} \phi_t\left(\sum_{n \in D_t} a_n z_n\right) &= \phi_t(a_{n_0} z_{n_0}) + \theta_k \sum_{n_0 < n \in I_t} \sum_{s \in J_n} \phi_s(a_n z_n) + \theta_k \sum_{s \in B} \phi_s\left(\sum_{n \in D_s} a_n z_n\right) \\ &\leq 2\phi_t(a_{n_0} z_{n_0}) + \theta_k \sum_{s \in B} C(f_{s,1} + f_{s,2})\left(\sum_{n \in D_s} a_n y_n\right) \\ &\leq C f_{t,1}(a_{n_0} y_{n_0}) + \theta_k \sum_{s \in B} C(f_{s,1} + f_{s,2})\left(\sum_{n \in D_s} a_n y_n\right) \\ &= C(f_{t,1} + f_{t,2})\left(\sum_{n \in D_t} a_n y_n\right). \end{aligned}$$

If the second term dominates the first one we repeat the argument by multiplying the estimation on $\phi_t(\sum_{n \in I_t} a_n z_n)$ by 2. $\square$

Remark 4.12. The reasoning presented in the proof of Theorem 4.10 can be adapted in case of $\theta = 1$ to obtain a version of Proposition 5.7 [17], giving equivalence of (y_n) of long "special" averages to some subsequence of the basis - it is enough to use the fact that in this case unit basis vectors (e_n) satisfies the assertion of Corollary 4.9.

Remark 4.13. Notice that as in the case of p -spaces the above proof provides in every block subspace of X some subsequentially minimal block subspace. Indeed any block sequence of vectors of the form $(y_1 + \dots + y_N)/\|y_1 + \dots + y_N\|$, where $y_1, \dots, y_N$ are chosen as in the beginning of the proof of Theorem 4.10 for arbitrary big N and $m_1, \dots, m_N$, spans a subsequentially minimal subspace.

We recall here that by Theorem 35 [13] if $0 < \inf c_n \leq \sup c_n < 1$ then the space $X = T[(\mathcal{S}_n, \theta_n)_n]$ fails to be a subsequentially minimal space in a strong way: X is saturated with block subspaces in which no block sequence is equivalent to a subsequence of the basis. It was also shown that the assumption $\sup_n c_n < 1$ is not necessary to have the property described above.

5. A NOTE FOR THE QUASIMINIMALITY OF THE DUAL SPACES

In this section we give a general result which allows to transfer the minimality and quasiminimality properties from a space to its dual space.

Theorem 5.1. *Let X be a Banach space with shrinking basis $(e_n)_{n \in \mathbb{N}}$ and let $(x_n^*)_{n \in \mathbb{N}}$ be a block sequence of $(e_n^*)_n$ such that there exist a block sequence $(x_n)_n$ of the basis such that*

- (1) $\frac{1}{1+\varepsilon} \leq \|x_n\| \leq 1 + \varepsilon$ and $x_n^*(x_n) = 1$.
- (2) $m \|\sum_n a_n e_{k_n}\| \leq \|\sum_n a_n x_n\| \leq M \|\sum_n a_n e_{k_n}\|$ for some $k_n \rightarrow \infty$.
- (3) *The map $P : X \rightarrow [x_n]$ defined by $P(x) = \sum_n x_n^*(x) x_n$ is a bounded projection.*

Then $(x_n^)_n$ is equivalent to the subsequence $(e_{k_n}^*)$.*

Proof. From the inequality $\|\sum_n a_n x_n\| \leq M \|\sum_n a_n e_{k_n}\|$ it follows that

$$\left\| \sum_n c_n e_{k_n}^* \right\| \leq M \left\| \sum_n c_n x_n^* \right\|$$

For the converse inequality let $x \in \mathcal{S}_X$ be such that $\sum_n c_n x_n^*(x) \geq (1+\varepsilon)^{-1} \|\sum_n c_n x_n^*\|$. We have that

$$\left\| \sum_n x_n^*(x) e_{k_n} \right\| \leq \left\| \sum_n x_n^*(x) x_n \right\| = \|P(x)\| \leq \|P\|$$

It follows that

$$\begin{aligned} \left\| \sum_n c_n e_{k_n}^* \right\| &\geq \|P\|^{-1} \sum_n c_n e_{k_n}^* \left(\sum_m x_m^*(x) e_{k_m} \right) \\ &= \|P\|^{-1} \sum_n c_n x_n^*(x) \geq ((1+\varepsilon)\|P\|)^{-1} \left\| \sum_n c_n x_n^* \right\| \end{aligned}$$

□

Corollary 5.2. *The dual space of Schlumprecht space $S = T[(\mathcal{A}_n, 1/\log_2(n+1))_n]$ is minimal ([9]). Dual spaces of mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ with $\sup \theta_n^{1/n} = 1$ are quasiminimal.*

Proof. In order to prove the Corollary we recall the following

Definition 5.3. *Let X be a Banach space with a basis. A normalized vector x is called a $(1+\varepsilon) - c_0^N$ -average, if $x = \sum_{k=1}^N x_k$ for some finite block sequence $(x_k)_{k=1}^N$ with $(1+\varepsilon)^{-1} \leq \|x_k\| \leq 1$ for all k .*

Remark 5.4. Assume X is a Banach space with a shrinking unconditional basis. With any $(1+\varepsilon) - c_0^N$ -average $x^* = \sum_{k=1}^N x_k^* \in X^*$ we can associate $(1+\varepsilon) - \ell_1^N$ -average in X in the following way: for every k pick a normalized $x_k \in X$ be such that $\text{supp } x_k = \text{supp } x_k^*$ and $x_k^*(x_k) = \|x_k^*\| \geq (1+\varepsilon)^{-1}$. Then the vector $x = \sum_{k=1}^N x_k / \left\| \sum_{k=1}^N x_k \right\|$ is $(1+\varepsilon) - \ell_1^N$ -average, since for any non-negative scalars $(a_k)_{k=1}^N$ we have

$$\left\| \sum_k a_k x_k \right\| \geq x^* \left(\sum_k a_k x_k \right) = \sum_{k=1}^N a_k x_k^*(x_k) \geq (1+\varepsilon)^{-1} \sum_k a_k.$$

Definition 5.5. [6] *Let X be a Banach space with the basis $(e_k)_{k \in \mathbb{N}}$. Let $\varepsilon > 0$ and $j \in \mathbb{N}$, $j > 1$. A convex combination $\sum_{k \in F} a_k e_k$ of the basis $(e_k)_{k \in \mathbb{N}}$ is called an $(\varepsilon; j)$ basic special convex combination if $F \in \mathcal{S}_j$ and $\sum_{k \in G} a_k < \varepsilon$ for every $G \in \mathcal{S}_{j-1}$.*

Let $(x_k)_{k \in \mathbb{N}}$ be a block sequence. A convex combination $\sum_{k \in F} a_k x_k$ of the sequence $(x_k)_{k \in \mathbb{N}}$ is called an $(\varepsilon; j)$ special convex combination $((\varepsilon; j)$ s.c.c.) of $(x_k)_{k \in \mathbb{N}}$ if $\sum_{k \in F} a_k e_{t_k}$ (where $t_k = \min \text{supp } x_k$ for each k) is an $(\varepsilon; j)$ basic special convex combination.

In order to apply Theorem 5.1 we need the following result.

Lemma 5.6. *Let $\varepsilon \in (0, 1)$ and Y^* be a subspace of S^* , the dual space to Schlumprecht space. Then for every $N \in \mathbb{N}$ there exists a $(1+\varepsilon) - c_0^N$ -average $z^* \in Y^*$.*

Let $\varepsilon \in (0, 1)$ and Y^ be a subspace of the dual space of a mixed Tsirelson space $X = T[(\mathcal{S}_n, \theta_n)_n]$ with $\sup \theta_n^{1/n} = 1$. Then for every $N \in \mathbb{N}$ there exists an (ε, N) s.c.c. $z = \sum_{i \in F} a_i z_i \in X$ and $z^* \in Y^*$ with $z^*(z) \geq 1 - \varepsilon$.*

The above Lemma for mixed Tsirelson spaces $T[(\mathcal{S}_n, \theta_n)_n]$ has been proved in [6], Proposition 5.4. For Schlumprecht space this lemma appeared in [9], Prop. 5.

We are now ready to complete the proof of the corollary.

Let (z_k^*) be a block sequence of $2 - c_0^{N_k}$ -averages in the block subspace Y^* of S^* and $(z_k)_k$ be the sequence of the associated $\ell_1^{N_k}$ -averages (Remark 5.4). We assume that the sequence N_k is fast increasing. By [3], see also [17], the sequence $(z_k)_k$ is equivalent to the basis of S and moreover spans a complemented subspace of S . Theorem 5.1 implies the result for the space S .

For the mixed Tsirelson spaces the analogous properties of a rapidly increasing sequence of s.c.c. averages have been proved for example in [17], thus again application of Theorem 5.1 finishes the proof of Corollary. □

Remark 5.7. The dual spaces of the spaces $T[\mathcal{S}_\alpha, \theta]$ are minimal. Indeed, we use the following result:

Proposition 5.8. [15] *Let normalized block sequences $(x_n), (y_n) \subset T[\mathcal{S}_\alpha, \theta]$ be such that $x_n < y_n < x_{n+1}$ ($n \in \mathbb{N}$). Then (x_n) and (y_n) are $24\theta^{-2}$ -equivalent.*

Standard dual argument yields that the same property holds also in the dual space i.e. if any normalized block sequences $(x_n^*)_n, (y_n^*)_n \subset T[\mathcal{S}_\alpha, \theta]$ with $x_n^* < y_n^* < x_{n+1}^*, n \in \mathbb{N}$ are equivalent. Now since c_0 is block finitely represented in the dual space $T[\mathcal{S}_\alpha, \theta]^*$ we follow the proof of [8] for Tsirelson space $T[\mathcal{S}_1, \theta]$ to obtain the minimality.

REFERENCES

- [1] D. Alspach and S.A. Argyros, *Complexity of weakly null sequences*, Diss. Math. **321**, 1–44 (1992).
- [2] G. Androulakis and E. Odell, *Distorting mixed Tsirelson spaces*, Israel J. Math. **109**, 125–149 (1999).
- [3] G. Androulakis and T. Schlumprecht, *The Banach space S is complementably minimal and subsequentially prime*, Studia Math. **156** no. 3, 227–242 (2003).
- [4] S.A. Argyros and I. Deliyanni, *Examples of asymptotic ℓ_1 Banach Spaces*, Trans. Amer. Math. Soc. **349**, 973–995 (1997).
- [5] S.A. Argyros, I. Deliyanni, and A. Manoussakis, *Distortion and spreading models in modified mixed Tsirelson spaces*, Studia Math. **157**, no. 3, 199–236 (2003).
- [6] S.A. Argyros and A. Tolias, *Methods in the theory of hereditarily indecomposable Banach spaces*, Mem. Amer. Math. Soc. **170**, no. 806 (2004).
- [7] J. Bernues and I. Deliyanni, *Families of finite subsets of $\mathbb{N}$ of low complexity and Tsirelson type spaces*, Math. Nachr. **222**, 15–29 (2001).
- [8] P. Casazza and E. Odell, *Tsirelson's space and minimal subspaces*, Texas functional analysis seminar 1982–1983 (Austin, Tex.), 61–72, Longhorn Notes, Univ. Texas Press, Austin, TX, 1983.
- [9] P. Casazza, N. Kalton, D. Kutzarova, and M. Mastyło, *Complex interpolation and complementably minimal spaces*. Interaction between functional analysis, harmonic analysis, and probability (Columbia, MO, 1994), 135–143, Lecture Notes in Pure and Appl. Math., **175**, Dekker, New York, 1996.
- [10] S. Dilworth, V. Ferenczi, D. Kutzarova, and E. Odell, *On strongly asymptotic ℓ_p spaces and minimality*, J. Lond. Math. Soc. (2) **75**, no. 2, 409–419 (2007).
- [11] V. Ferenczi and C. Rosendal, *Banach spaces without minimal subspaces*, to appear in J. Funct. Anal.
- [12] M. Junge, D. Kutzarova, and E. Odell, *On asymptotically symmetrical Banach spaces*, Studia Math. **173**, no. 3, 203–231 (2006).
- [13] D. Kutzarova, D. Leung, A. Manoussakis, and W.-K. Tang, *Minimality properties of Tsirelson type spaces*, Studia Math. **187**, no. 3, 233–263 (2008).
- [14] D. Leung and W.-K. Tang, *ℓ_1 -spreading models in subspaces of mixed Tsirelson spaces*, Studia Math. **172**, no. 1, 47–128 (2006).
- [15] J. Lopez-Abad and A. Manoussakis, *A Classification of Tsirelson Type Spaces*, Canadian Journal of Math. **60**, no.5, 1108–1048 (2008).
- [16] A. Manoussakis, *A note on certain equivalent norms on Tsirelson's space*, Glasgow Math. J. **46**, 379–390 (2004).
- [17] A. Manoussakis, *On the structure of a certain class of mixed Tsirelson spaces*, Positivity **5**, no. 3, 193–238 (2001).
- [18] B. Maurey, *Type, cotype and K -convexity*, Handbook of the geometry of Banach spaces, Vol. 2, 1299–1332, North-Holland, Amsterdam, 2003.
- [19] E. Odell and N. Tomczak-Jaegermann, *On certain equivalent norms on Tsirelson's space*, Illinois J. Math. **44**, 51–71 (2000).
- [20] E. Odell, N. Tomczak-Jaegermann, and R. Wagner, *Proximity to ℓ_1 and Distortion in Asymptotic ℓ_1 Spaces*, J. Funct. Anal. **150**, 101–145 (1997).
- [21] T. Schlumprecht, *An arbitrary distortable Banach space*, Israel J. Math. **76**, 81–95 (1991).
- [22] T. Schlumprecht, *How many operators exist on a Banach space?* Trends in Banach spaces and operator theory (Memphis, TN, 2001), 295–333, Contemp. Math. **321**, Amer. Math. Soc., Providence, RI, 2003.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AEGEAN, KARLOVASI, SAMOS, GR 83200,
GREECE

E-mail address: amanouss@aegean.gr

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW,
POLAND

E-mail address: anna.pelczar@im.uj.edu.pl